


No.

200100014



THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

The J. C. Robinson Seed Company

Whereas, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR IMPORTING IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE PURPOSE, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

CORN, FIELD

'JCR503'

In Testimony Whereof, I have hereunto set my hand and caused the seal of the Plant Variety Protection Office to be affixed at the City of Washington, D.C. this tenth day of April, in the year two thousand three.

Attest:




[Signature]

Commissioner
Plant Variety Protection Office
Agricultural Marketing Service

[Signature]

Secretary of Agriculture

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY - PLANT VARIETY PROTECTION OFFICE APPLICATION FOR PLANT VARIETY PROTECTION CERTIFICATE <i>(Instructions and information collection burden statement on reverse)</i>		The following statements are made in accordance with the Privacy Act of 1974 (5 U.S.C. 552a) and the Paperwork Reduction Act (PRA) of 1995. Application is required in order to determine if a plant variety protection certificate is to be issued (u U.S.C. 2421). Information is held confidential until certificate is issued (7 U.S.C. 2426).	
1. NAME OF OWNER The J. C. Robinson Seed Company		2. TEMPORARY DESIGNATION OR EXPERIMENTAL NAME	3. VARIETY NAME JCR503
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP Code, and Country) 100 J. C. Robinson Blvd PO Box A Waterloo, Ne 68069		5. TELEPHONE (include area code) 402-289-0252	FOR OFFICIAL USE ONLY PVPO NUMBER 200100014
		6. FAX (include area code) 402-779-4370	FILING DATE Oct. 18, 2000
7. IF THE OWNER NAMED IS NOT A "PERSON", GIVE FORM OF ORGANIZATION (corporation, partnership, association, etc.) Corporation	8. IF INCORPORATED, GIVE STATE OF INCORPORATION Nebraska	9. DATE OF INCORPORATION 8-1-1964	
10. NAME AND ADDRESS OF OWNER REPRESENTATIVE(S) TO SERVE IN THIS APPLICATION. (First person listed will receive all papers) Warren Anthony Foundation Seed Manager 100 J. C. Robinson Blvd PO Box A Waterloo, Ne 68069		F E E S R E C E I V E D	FILING AND EXAMINATION FEES: \$ 2450 + 255
			DATE 8/24/00 + 10/18/00 CERTIFICATION FEE: \$ 432.00 DATE 3/24/00
11. TELEPHONE (include area code) 402-289-0252	12. FAX (include area code) 402-779-4307	13. E_MAIL wanthony@jcrob.com	14. CROP KIND (Common Name) Corn
15. GENUS AND SPECIES NAME OF CROP Zae Mays E.		16. FAMILY NAME (Botanical) Granineae	17. IS THE VARIETY A FIRST GENERATION HYBRID? <input type="checkbox"/> YES <input checked="" type="checkbox"/> NO
18. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED (Follow instructions on reverse) a. <input checked="" type="checkbox"/> Exhibit A. Origin and Breeding History of the Variety b. <input checked="" type="checkbox"/> Exhibit B. Statement of Distinctness c. <input checked="" type="checkbox"/> Exhibit C. Objective Description of Variety d. <input checked="" type="checkbox"/> Exhibit D. Additional Description of the Variety (Optional) e. <input checked="" type="checkbox"/> Exhibit E. Statement of the Basis of the Owner's Ownership f. <input checked="" type="checkbox"/> Voucher Sample (2,500 viable untreated seeds or, for tuber propagated varieties, verification that tissue culture will be deposited and maintained in an approved public repository) g. <input checked="" type="checkbox"/> Filing and Examination Fee (\$2,450), made payable to "Treasurer of the United States" (Mail to the Plant Variety Protection Office)		19. DOES THE OWNER SPECIFY THAT SEED OF THIS VARIETY BE SOLD AS A CLASS OF CERTIFIED SEED? (See Section 83(a) of the Plant Variety Protection Act) <input type="checkbox"/> YES (If "yes", answer items 20 and 21 below) <input type="checkbox"/> NO (if "no," go to item 22)	
		20. DOES THE OWNER SPECIFY THAT SEED OF THIS VARIETY BE LIMITED AS TO NUMBER OF CLASSES? <input type="checkbox"/> YES <input checked="" type="checkbox"/> NO IF YES, WHICH CLASSES? <input type="checkbox"/> FOUNDATION <input type="checkbox"/> REGISTERED <input type="checkbox"/> CERTIFIED	
		21. DOES THE OWNER SPECIFY THAT THE CLASSES BE LIMITED AS TO THE NUMBER OF GENERATIONS? <input type="checkbox"/> YES <input checked="" type="checkbox"/> NO IF YES, SPECIFY THE <input type="checkbox"/> FOUNDATION <input type="checkbox"/> REGISTERED <input type="checkbox"/> CERTIFIED NUMBER 1,2,3, etc. (If additional explanation is necessary, please use the space indicated on the reverse.)	
22. HAS THE VARIETY (INCLUDING ANY HARVESTED MATERIAL) OR A HYBRID PRODUCED FROM THIS VARIETY BEEN SOLD, DISPOSED OF, TRANSFERRED, OR USED IN THE U.S. OR OTHER COUNTRIES? <input type="checkbox"/> YES <input checked="" type="checkbox"/> NO IF YES, YOU MUST PROVIDE THE DATE OF FIRST SALE, DISPOSITION, TRANSFER, OR USE FOR EACH COUNTRY AND THE CIRCUMSTANCES (Please use space indicated on reverse.)		23. IS THE VARIETY OR ANY COMPONENT OF THE VARIETY PROTECTED BY INTELLECTUAL PROPERTY RIGHT (PLANT BREEDER'S RIGHT OR PATENT)? <input type="checkbox"/> YES <input checked="" type="checkbox"/> NO IF YES, GIVE COUNTRY, DATE OF FILING OR ISSUANCE AND ASSIGNED REFERENCE NUMBER. (Please use space indicated on reverse.)	
24. The owners declare that a viable sample of basic seed of the variety will be furnished with application and will be replenished upon request in accordance with such regulations as may be applicable, or for a tuber propagated variety a tissue culture will be deposited in a public repository and maintained for the duration of the certificate. The undersigned owner(s) is(are) the owner of this sexually reproduced or tuber propagated plant variety, and believe(s) that the variety is new, distinct, uniform, and stable as required in Section 42, and is entitled to protection under the provisions of Section 42 of the Plant Variety Protection Act. Owner(s) is(are) informed that false representation herein can jeopardize protection and result in penalties.			
SIGNATURE OF OWNER NAME (Please print or type) Warren Anthony		SIGNATURE OF OWNER  NAME (Please print or type) Warren Anthony	
CAPACITY OR TITLE DATE Foundation Seed cManager		CAPACITY OR TITLE DATE 03 Feb. 2003	

INSTRUCTIONS

GENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in the PVPO: (1) Completed application form signed by the owner; (2) completed exhibits A, B, C, E; (3) for a seed reproduced variety at least 2,500 viable untreated seeds, for a hybrid variety at least 2,500 untreated seeds of each line necessary to reproduce the variety, or for tuber reproduced varieties verification that a viable (in the sense that it will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; (4) check drawn on a U.S. bank for \$2,450 (\$300 filing fee and \$2,150 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice.) Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfiled. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 500, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initialed and dated. DO NOT use masking materials to make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$300 for issuance of the certificate. Certificates will be issued to owner, not licensee or agent

Plant Variety Protection Office

Telephone: (301)504-5518

Fax: (301)504-5291

Homepage: http://www.ams.usda.gov/science/pvp.htm

ITEM

- 18a. Give: (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method; (2) the details of subsequent stages of selection and multiplication; (3) evidence of uniformity and stability; and (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 18b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all difference objectively;
 - (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and
 - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 18c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- 18d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 18e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 19. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- 20. See Section 83 of the Act for the Contents and Term of Plant Variety Protection.
- 21. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 22. See Section 5.5 of the Act for instructions on claiming the benefit of earlier filing date.

21. CONTINUED FROM FRONT (Please provide a statement as to the limitation and sequence of generations that may be certified.)
No Limit

22. CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety (including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)

23. CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right, or Patent).)

USA only

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. There is no charge for filing a change of address. The fee for filing a change of ownership or assignment or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

To avoid conflict with other variety names in use, the applicant must check the variety names proposed by contacting: Seed Branch, AMS, USDA, Room 213, Building 306, Beltsville Agricultural Research Center—East, Beltsville, MD 20705. Telephone: (301)504-8089.

Public reporting burden for this collection of information is estimated to average 30 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Agriculture, Clearance Officer, OIRM, AG Box 7630, Jamie L. Whitten Building, Washington, D.C. 20250. When replying, refer to OMB No. 0581-0055 and form number in your letter. Under the PRA of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number

The U. S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and marital or familial status. (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact USDA Office of Communications at (202)720-2791. To file a complaint, write secretary of Agriculture, Washington D.C. 20250, or call (202)720-7327 (voice) or (202)720-1127 (TDD). USDA is an equal opportunity employer.

Origin and Breeding History of the Inbred JCR503

Exhibit A:

JCR503 is a corn inbred line developed from the single cross of LH195 x SR108 using the pedigree method of breeding. Selfing and selection were conducted for six generations in the development of JCR503. The selection criteria used in the development of JCR503 were: grain yield; high plant density tolerance; stalk and root strength; late plant greenness; and disease tolerance. Testcrosses with unrelated inbreds were made and evaluated over multiple years and locations in the promotion of JCR503 to commercial status.

LH195, a progenitor of JCR503, is a proprietary field inbred line developed by Holden's Foundation Seeds, Inc. Holden's was granted plant variety protection certificate #9000047 for inbred LH195. SR108, a progenitor of JCR503, is a proprietary field corn inbred of The J.C. Robinson Seed Company. SR108 was developed from a proprietary synthetic population containing LH132 and an exotic germplasm source containing B73 and Tuxpeno.

JCR503 has shown uniformity and stability for all traits as described in Exhibit C – "Objective Description of Variety". It has been self-pollinated and ear-rowed a sufficient number of generations, with careful attention given to uniformity of plant type to ensure homozygosity and phenotypic purity. Several S4 ear-rows were grown at Henrietta, MO during the summer of 1996. A uniform S4 was selected and several S5 progeny were then grown in Molokai, HA during the winter of 1996-1997. A uniform S5 was selected and several S6 progeny were then grown in Elk City, NE during the summer of 1997. These ear rows were evaluated to assure they were uniform and identical for phenotype. The S7 seed harvested from Elk City was bulked and grown in hand-pollinated increases in both Ponce, Puerto Rico and Molokai, HA during the winter of 1997-1998. The hand-pollinated increases were observed by the breeder and assistant breeder to assure uniformity of the inbred line. The S8 seed harvested from Hawaii and Puerto Rico was grown in an open-pollinated line increase isolation at Waterloo, NE during the summer of 1998. The J.C. Robinson foundation seed manager and breeder of JCR503 both observed this increase and confirmed the uniformity of the inbred line. A subsequent open-pollinated increase grown in 1999 was also found to be uniform.

No variant traits have been observed or are expected in JCR503.

Developmental History for JCR503:

<u>Location/Season/ Year</u> :	<u>Inbreeding Level:</u>
Molokai, HA/Winter/1993-1994	S0 self
Henrietta, MO/Summer/1994	S1 self and select
Henrietta, MO/Summer/1995	S2 self and select
Molokai, HA/Winter/1995-96	S3 self
Henrietta, MO/Summer/1996	S4 self and select
Ponce, Puerto Rico/Winter/1996-97	S5 self
Elk City, NE/Summer/1997	S6 self and initial bulk
Ponce, Puerto Rico & Molokai, HA /Winter/1997-1998	S7 self and hand increase of bulk
Waterloo, NE/Summer/1998	open-pollinated foundation increase
Waterloo, NE/Summer/1999	open-pollinated foundation increase

Statement of Distinctness for Inbred JCR503

Exhibit B:

JCR503 is most similar to B73, however, there are distinguishing differences between the two inbreds. The most distinctive differences between the lines are those of silk color and cob color. Silks of JCR503 are pink colored whereas those of B73 are green-yellow. When using the Munsell Color Charts for Plant Tissues as a reference, the silk color of JCR503 would be classified as $7.5YR7/6$ and that of B73 would be $5R6/6$. The cob color of JCR503 is white (2.5Y8/6) and that of B73 is red (10R4/10).

The following traits are highly significant at the 1% level (Student's t-Test procedure).

Trait	JCR503		B73		Mean Diff	t-Value	Prob
	N	Mean	N	Mean			
Plant Height	15	243.1	15	228.7	14.4	5.66	0.0000
Length of Top Ear Internode	15	12.8	15	13.9	-1.1	-3.64	0.0011
Leaf Length	15	84.1	15	77.2	6.9	5.74	0.0000
Leaf Angle	15	12.7	15	13.5	-2.9	-3.29	0.0030
Tassel Length	15	48.5	15	38.6	9.9	10.33	0.0000
Ear Length	15	14.6	15	13.4	1.2	3.54	0.0014
Ear Diameter	15	45.5	15	41.6	3.9	5.26	0.0000
Ear Weight	15	136.8	15	82.3	54.5	6.55	0.0000
Shank Length	15	8.2	15	6.0	2.2	4.19	0.0003
Kernel Length	15	11.7	15	10.5	1.2	4.95	0.0000
Kernel Width	15	7.9	15	7.4	0.5	3.23	0.0030
Kernel Weight	15	32.2	15	26.6	5.6	6.33	0.0000

United States Department of Agriculture, Agricultural Marketing Service
Science Division, Plant Variety Protection Office
National Agricultural Library Building, Room 500
Beltsville, MD 20705

OBJECTIVE DESCRIPTION OF VARIETY
CORN (*Zea mays L.*)

Name of Applicant(s) The J. C. Robinson Seed Company	Variety Seed Source	Variety Name or Temporary Designation JCR503
Address (Street & No., or R.F.D. No., City, State, Zip Code and Country) 100 J. C. Robinson Blvd PO Box A Waterloo, Ne 68069		FOR OFFICIAL USE PVPO Number 200100014

Place the appropriate number that describes the varietal characters typical of this inbred variety in the spaces below. Right justify whole numbers by adding leading zeroes if necessary. Completeness should be striven for to establish an adequate variety description. Traits designated by a "*" are considered necessary for an adequate variety description and must be completed.

COLOR CHOICES (Use in conjunction with Munsell color code to describe all color choices: describe #25 and #26 in Comments section):

01 = Light Green	06 = Pale Yellow	11 = Pink	16 = Pale Purple	21 = Buff
02 = Medium Green	07 = Yellow	12 = Light Red	17 = Purple	22 = Tan
03 = Dark Green	08 = Yellow-Orange	13 = Cherry Red	18 = Colorless	23 = Brown
04 = Very Dark Green	09 = Salmon	14 = Red	19 = White	24 = Bronze
05 = Green-Yellow	10 = Pink-Orange	15 = Red & White	20 = White Capped	25 = Variegated (Describe)
				26 = Other (Describe)

STANDARD INBRED CHOICES (Use the most similar (in background and maturity) of these to make comparisons based on grow-out trial data):

Yellow Dent Families:		Yellow Dent (Unrelated):		Sweet Corn:
Family	Members	Co109	ND246	C13, Iowa5125, P39, 2132
B14	CM105, A632, B64, B68	Oh7,	T232	
B37	B37, B76, H84	W117	W153R	Popcorn:
B73	N192, A679, B73, NC268	W182BN		SG1533, 4722, HP301, HP7211
C103	Mo17, Va102, Va35, A682			
Oh43	A619, MS71, H99, Va26	White Dent:		Pipecorn:
WF9	W64A, A554, A654, Pa91	Cl66, H105, Ky228		Mo15W, Mo16W, Mo24W

1. TYPE: (describe intermediate types in Comments Section) * <u>2</u> 1=Sweet 2=Dent 3=Flint 4=Flour 5=Pop 6=Ornamental 7=Pipecorn	Standard Inbred Name B73 <u>2</u>																																										
2. REGION WHERE DEVELOPED IN THE U.S.A.: * <u>2</u> 1=Northwest 2=Northcentral 3=Northeast 4=Southeast 5=Southcentral 6=Southwest 7=Other	Standard Seed Source _____ <u>2</u>																																										
3. MATURITY (In Region Best Adaptability: show Heat unit formula in "Comments" section): <table border="0"> <tr> <td>DAYS</td> <td>HEAT UNITS</td> <td></td> <td>DAYS</td> <td>HEAT UNITS</td> </tr> <tr> <td>* <u>067</u></td> <td><u>1312.5</u></td> <td>From emergence to 50% of plants in silk</td> <td><u>065</u></td> <td><u>1254.0</u></td> </tr> <tr> <td>* <u>067</u></td> <td><u>1312.5</u></td> <td>From emergence to 50% of plants in pollen</td> <td><u>064</u></td> <td><u>1224.0</u></td> </tr> <tr> <td><u>003</u></td> <td><u>0090.0</u></td> <td>From 10% to 90% pollen shed</td> <td><u>002</u></td> <td><u>0060.5</u></td> </tr> <tr> <td>* ---</td> <td>-----</td> <td>From 50% silk to optimum edible quality</td> <td>---</td> <td>-----</td> </tr> <tr> <td>---</td> <td>-----</td> <td>From 50% silk to harvest at 25% moisture</td> <td>---</td> <td>-----</td> </tr> </table>	DAYS	HEAT UNITS		DAYS	HEAT UNITS	* <u>067</u>	<u>1312.5</u>	From emergence to 50% of plants in silk	<u>065</u>	<u>1254.0</u>	* <u>067</u>	<u>1312.5</u>	From emergence to 50% of plants in pollen	<u>064</u>	<u>1224.0</u>	<u>003</u>	<u>0090.0</u>	From 10% to 90% pollen shed	<u>002</u>	<u>0060.5</u>	* ---	-----	From 50% silk to optimum edible quality	---	-----	---	-----	From 50% silk to harvest at 25% moisture	---	-----													
DAYS	HEAT UNITS		DAYS	HEAT UNITS																																							
* <u>067</u>	<u>1312.5</u>	From emergence to 50% of plants in silk	<u>065</u>	<u>1254.0</u>																																							
* <u>067</u>	<u>1312.5</u>	From emergence to 50% of plants in pollen	<u>064</u>	<u>1224.0</u>																																							
<u>003</u>	<u>0090.0</u>	From 10% to 90% pollen shed	<u>002</u>	<u>0060.5</u>																																							
* ---	-----	From 50% silk to optimum edible quality	---	-----																																							
---	-----	From 50% silk to harvest at 25% moisture	---	-----																																							
4. PLANT: <table border="0"> <tr> <td></td> <td>Standard Deviation</td> <td>Sample Size</td> <td></td> <td>Standard Deviation</td> <td>Sample Size</td> </tr> <tr> <td>* <u>243.1</u></td> <td>cm Plant Height (to tassel tip)</td> <td><u>4.9</u></td> <td><u>15</u></td> <td><u>228.7</u></td> <td><u>8.4</u></td> </tr> <tr> <td>* <u>102.9</u></td> <td>cm Ear Height (to base of top ear node)</td> <td><u>7.5</u></td> <td><u>15</u></td> <td><u>108.9</u></td> <td><u>4.7</u></td> </tr> <tr> <td><u>012.8</u></td> <td>Cm Length of Top Ear Internode</td> <td><u>0.67</u></td> <td><u>15</u></td> <td><u>013.9</u></td> <td><u>0.9</u></td> </tr> <tr> <td><u>0.0</u></td> <td>Average Number of Tillers</td> <td><u>0</u></td> <td><u>15</u></td> <td><u>000.0</u></td> <td><u>0</u></td> </tr> <tr> <td>* <u>1.0</u></td> <td>Average Number of Ears per Stalk</td> <td><u>0</u></td> <td><u>15</u></td> <td><u>001.0</u></td> <td><u>0</u></td> </tr> <tr> <td><u>2</u></td> <td>Anthocyanin of Brace Roots: 1=Absent 2=Faint 3=Moderate 4=Dark</td> <td></td> <td></td> <td><u>4</u></td> <td></td> </tr> </table>		Standard Deviation	Sample Size		Standard Deviation	Sample Size	* <u>243.1</u>	cm Plant Height (to tassel tip)	<u>4.9</u>	<u>15</u>	<u>228.7</u>	<u>8.4</u>	* <u>102.9</u>	cm Ear Height (to base of top ear node)	<u>7.5</u>	<u>15</u>	<u>108.9</u>	<u>4.7</u>	<u>012.8</u>	Cm Length of Top Ear Internode	<u>0.67</u>	<u>15</u>	<u>013.9</u>	<u>0.9</u>	<u>0.0</u>	Average Number of Tillers	<u>0</u>	<u>15</u>	<u>000.0</u>	<u>0</u>	* <u>1.0</u>	Average Number of Ears per Stalk	<u>0</u>	<u>15</u>	<u>001.0</u>	<u>0</u>	<u>2</u>	Anthocyanin of Brace Roots: 1=Absent 2=Faint 3=Moderate 4=Dark			<u>4</u>		
	Standard Deviation	Sample Size		Standard Deviation	Sample Size																																						
* <u>243.1</u>	cm Plant Height (to tassel tip)	<u>4.9</u>	<u>15</u>	<u>228.7</u>	<u>8.4</u>																																						
* <u>102.9</u>	cm Ear Height (to base of top ear node)	<u>7.5</u>	<u>15</u>	<u>108.9</u>	<u>4.7</u>																																						
<u>012.8</u>	Cm Length of Top Ear Internode	<u>0.67</u>	<u>15</u>	<u>013.9</u>	<u>0.9</u>																																						
<u>0.0</u>	Average Number of Tillers	<u>0</u>	<u>15</u>	<u>000.0</u>	<u>0</u>																																						
* <u>1.0</u>	Average Number of Ears per Stalk	<u>0</u>	<u>15</u>	<u>001.0</u>	<u>0</u>																																						
<u>2</u>	Anthocyanin of Brace Roots: 1=Absent 2=Faint 3=Moderate 4=Dark			<u>4</u>																																							

5

Application Variety Data			Page 2	Standard Inbred Data		
5. Leaf:			Standard Deviation	Sample Size	Standard Deviation Sample Size	
*	<u>011.0</u>	cm Width of Ear Node Leaf	<u>1.03</u>	<u>15</u>	<u>010.6</u>	<u>0.43</u> <u>15</u>
*	<u>084.1</u>	cm Length of Ear Node Leaf	<u>3.57</u>	<u>15</u>	<u>077.2</u>	<u>2.98</u> <u>15</u>
*	<u>06</u>	Number of leaves above top ear	<u>0.61</u>	<u>15</u>	<u>05</u>	<u>0.45</u> <u>15</u>
	<u>013</u>	Degrees Leaf Angle (measure from 2 nd leaf above ear at anthesis to stalk above leaf)	<u>2.43</u>	<u>15</u>	<u>016</u>	<u>2.32</u> <u>15</u>
*	<u>03</u>	Leaf Color (Munsell code <u>5GY4/4</u>)			<u>04</u>	(Munsell code <u>5GY3/4</u>)
	<u>2</u>	Leaf Sheath Pubescence (Rate on scale from 1=none to 9=like peach fuzz)			<u>8</u>	
	<u>4</u>	Marginal Waves (Rate on scale from 1=none to 9=many)			<u>4</u>	
	<u>6</u>	Longitudinal Creases (Rate on scale from 1=none to 9=many)			<u>5</u>	
6. Tassel:			Standard Deviation	Sample Size	Standard Deviation Sample Size	
*	<u>11</u>	Number of primary Lateral Branches	<u>2.35</u>	<u>15</u>	<u>09</u>	<u>2.02</u> <u>15</u>
	<u>029</u>	Branch Angle from Central Spike	<u>4.11</u>	<u>15</u>	<u>021</u>	<u>8.6</u> <u>15</u>
*	<u>48.5</u>	cm Tassel Length (from top leaf collar to tassel tip)	<u>2.56</u>	<u>15</u>	<u>38.6</u>	<u>2.66</u> <u>15</u>
	<u>5</u>	Pollen Shed (Rate on scale from 0=male sterile to 9=heavy shed)			<u>4</u>	
	<u>07</u>	Anther Color (Munsell code <u>5Y8/8</u>)			<u>07</u>	(Munsell code <u>5Y8/8</u>)
	<u>05</u>	Glume Color (Munsell code <u>2.5GY8/8</u>)			<u>05</u>	(Munsell code <u>2.5GY8/6</u>)
	<u>1</u>	Bar Glumes (Glume Bands) 1 = Absent 2 = Present			<u>1</u>	
7a. EAR (Unhusked data):						
*	<u>11</u>	Silk Color (3 days after emergence) (Munsell code 7.5YR7/6 ^{5R 6/6})			<u>05</u>	(Munsell code <u>2.5GY8/8</u>)
	<u>05</u>	Fresh Husk Color (25 days after 50% silking)(Munsell code <u>2.5GY7/6</u>)			<u>02</u>	(Munsell code <u>5GY6/6</u>)
	<u>21</u>	Dry Husk Color (65 days after 50% silking) (Munsell code <u>5Y8/8</u>)			<u>21</u>	(Munsell code <u>5Y8/8</u>)
*	<u>1</u>	Position of Ear at Dry Husk Stage: 1=Upright 2=Horizontal 3=Pendent			<u>1</u>	
	<u>4</u>	Husk Tightness (Rate on scale from 1=Very loose to 9=Very tight)			<u>6</u>	
	<u>3</u>	Husk Extension (at harvest): 1=Short (ears exposed) 2=Medium (<8 cm) 3=Long (8-10 cm beyond ear tip) 4=Very Long (>10 cm)			<u>1</u>	
7b. EAR (Husked Ear Data):			Standard Deviation	Sample Size	Standard Deviation Sample Size	
*	<u>14.6</u>	cm Ear Length	<u>1.13</u>	<u>15</u>	<u>13.4</u>	<u>0.67</u> <u>15</u>
*	<u>45.5</u>	mm Ear Diameter at mid-point	<u>2.36</u>	<u>15</u>	<u>41.6</u>	<u>1.63</u> <u>15</u>
	<u>136.8</u>	Gm Ear Weight	<u>24.7</u>	<u>15</u>	<u>082.3</u>	<u>20.5</u> <u>15</u>
*	<u>15</u>	Number of Kernel Rows	<u>1.98</u>	<u>15</u>	<u>16</u>	<u>1.03</u> <u>15</u>
	<u>2</u>	Kernel Rows: 1=Indistinct 2=Distinct			<u>2</u>	
	<u>1</u>	Row Alignment: 1=Straight 2=Slightly Curved 3=Spiral			<u>1</u>	
	<u>08.2</u>	cm Shank Length	<u>1.55</u>	<u>15</u>	<u>06.0</u>	<u>1.26</u> <u>15</u>
	<u>2</u>	Ear Taper: 1=Slight 2=Average 3=Extreme			<u>2</u>	
Application Variety Data			Standard Inbred Data			

5MS 5/20/02

NOTE: Use chart on first page to choose color codes for traits

Application Variety Data			Page 3	Standard Inbred Data		
8. KERNEL (Dried):			Standard Deviation	Sample Size	Standard Deviation	Sample Size
<u>1 1.7</u>	mm Kernel Length	<u>0.69</u>	<u>15</u>	<u>1 0.5</u>	<u>0.59</u>	<u>15</u>
<u>0 7.9</u>	mm Kernel Width	<u>0.51</u>	<u>15</u>	<u>0 7.4</u>	<u>0.24</u>	<u>15</u>
<u>0 4.8</u>	mm Kernel Thickness	<u>0.23</u>	<u>15</u>	<u>0 4.9</u>	<u>0.73</u>	<u>15</u>
<u>4 2.9</u>	% Round Kernels (Shape Grade)	<u>14.0</u>	<u>15</u>	<u>5 3.6</u>	<u>21.6</u>	<u>13</u>
<u>1</u>	Aleurone Color Pattern: 1=Homozygous 2=Segregating _____			<u>1</u>		
* <u>1 8</u>	Aleurone Color (Munsell code COLORLESS)			<u>1 8</u>	(Munsell code COLORLESS)	
* <u>0 7</u>	Hard Endosperm Color (Munsell code 2.5Y7/8)			<u>0 7</u>	(Munsell code 2.5Y6/8)	
* <u>0 3</u>	Endosperm Type: 1=Sweet (sul) 2=Extra Sweet (sh2) 3=Normal Starch 4=High Amylose Starch 5=Waxy Starch 6=High Protein 7= High Lysine 8=Super Sweet (se) 9=High Oil 10=Other _____			<u>0 3</u>		
<u>3 2.2</u>	gm Weight per 100 Kernels (unsized sample)	<u>3.09</u>	<u>15</u>	<u>2 6.6</u>	<u>1.70</u>	<u>13</u>
9. COB:			Standard Deviation	Sample Size	Standard Deviation	Sample Size
* <u>2 8.5</u>	mm Cob Diameter at mid-point	<u>1.77</u>	<u>15</u>	<u>2 7.8</u>	<u>1.18</u>	<u>15</u>
<u>1 9</u>	Cob Color (Munsell code 2.5Y8/6)			<u>1 4</u>	(Munsell code 10R4/10)	
10. DISEASE RESISTANCE (Rate from 1 (most susceptible) to 9 (most resistant): Leave blank if not tested: leave Race or Strain Options blank if polygenic):						
A. Leaf Blights, Wilts, and Local Infection Diseases						
_ Anthracnose Leaf Blight (<i>Colletotrichum graminicola</i>)			-			
_ Common Rust (<i>Puccinia sorghi</i>)			-			
_ Common Smut (<i>Ustilago maydis</i>)			-			
_ Eyespot (<i>Kabatiella zeae</i>)			-			
_ Goss's Wilt (<i>Clavibacter michiganense spp. nebraskense</i>)			-			
_ Gray Leaf Spot (<i>Cercospora zeae-maydis</i>)			-			
_ Helminthosporium Leaf Spot (<i>Bipolaris zeicola</i>) Race _____			_ Race _____			
_ Northern Leaf Blight (<i>Exserohilum turcicum</i>) Race _____			_ Race _____			
_ Southern Leaf Blight (<i>Bipolaris maydis</i>) Race _____			_ Race _____			
_ Southern Rust (<i>Puccinia polysora</i>)			-			
_ Stewart's Wilt (<i>Erwinia stewartii</i>)			-			
_ Other (Specify) _____			- _____			
B. Systemic Diseases						
_ Corn Lethal Necrosis (MCMV and MDMV)			-			
_ Head Smut (<i>Sphacelotheca reiliana</i>)			-			
_ Maize Chlorotic Dwarf Virus(MCDV)			-			
_ Maize Chlorotic Mottle Virus (MCMV)			-			
_ Maize Dwarf Mosaic Virus (MDMV) Strain _____			_ Strain _____			
_ Sorghum Downy Mildew of Corn (<i>Peronosclerospora sorghi</i>)			-			
_ Other (Specify) _____			- _____			
C. Stalk Rots						
_ Anthracnose Stalk Rot (<i>Colletotrichum graminicola</i>)			-			
_ Diplodia Stalk Rot (<i>Stenocarpella maydis</i>)			-			
_ Fusarium Stalk Rot (<i>Fusarium moniliforme</i>)			-			
_ Gibberella Stalk Rot (<i>Gibberella zeae</i>)			-			
_ Other (Specify) _____			- _____			
D. Ear and Kernel Rots						
_ Aspergillus Ear and Kernel Rot (<i>Aspergillus flavus</i>)			-			
_ Diplodia Ear Rot (<i>Stenocarpella maydis</i>)			-			
_ Fusarium Ear and Kernel Rot (<i>Fusarium moniliforme</i>)			-			
_ Gibberella Ear Rot (<i>Gibberella zeae</i>)			-			
_ Other (Specify) _____			- _____			
Application Variety Data			Standard Inbred Data			

NOTE: Use Chart on first page to choose color codes for color traits

11. INSECT RESISTANCE (Rate from 1 (most susceptible) to 9 (most resistant))

leave blank if not tested:

200100014

Standard Deviation Sample Size

- Banks Grass Mite (*Oligonychus pratensis*) Standard Deviation Sample Size
- Corn Earworm (*Helicoverpa zea*)
- Leaf-feeding
- Silk feeding:
- ___ mg Larval wt. _____
- Ear Damage
- Corn Leaf Aphid (*Rhopalosiphum maidis*)
- Corn Sap Beetle (*Carpophilus dimidiatus*)
- European Corn Borer (*Ostrinia nubilalis*)
- 1st Generation (Typically Whorl Leaf Feeding)
- 2nd Generation (Typically Leaf Sheath-Collar Feeding)
- Stalk Tunneling:
- ___ cm tunneled plant _____
- Fall Armyworm (*Spodoptera frugiperda*)
- Leaf Feeding
- Silk Feeding:
- ___ mg larval wt. _____
- Maize Weevil (*sitophilus zeamaize*)
- Northern Rootworm (*Diabrotica barberi*)
- Southern Rootworm (*Diabrotica undecimpunctata*)
- Southwestern Corn Borer (*Diatraea grandiosella*)
- Leaf Feeding
- Stalk Tunneling:
- ___ cm tunneled/plant _____
- Two-spotted Spider Mite (*Tetranychus urticae*)
- Western Rootworm (*Diabrotica virgifera virgifera*)
- Other (Specify) _____

12. AGRONOMIC TRAITS:

- Stay Green (at 65 days after anthesis) (Rate on a scale from 1=Worst to 9=Excellent.)
- ___ %Dropped Ears (at 65 day anthesis)
- ___ %Pre-anthesis Brittle Snapping
- ___ %Pre-anthesis Root Lodging
- ___ %Post-anthesis Root Lodging (at 65 days after anthesis)
- ___ Kg/ha Yield of Inbred Per Se (at 12-13% grain moisture)

13. MOLECULAR MARKERS: (0=data unavailable 1=data available but not supplied 2=data supplied)

JMS 5/20/02

- Isozymes RFLP's RAPD's

REFERENCES:

Butler, D.R. 1954. A System for the Classification of Corn Inbred Lines. PhD Thesis, Ohio State University.
 Emerson, R.A., G.W. Beadle, and A.C. Fraser. 1935. A Summary of Linkage Studies in Maize. Cornell A.E.S., Mem. 180.
 Farr, D.F., G.F. Bills, G.P. Chamuris, A.Y. Rossman, 1989. Fungi on Plant and Plant Products in the United States. The AmericanPhytopathological Society, St. Paul, MN.
 Inglett, G.E. (Ed.) 1970. Corn: Culture, Processing, Products. Avi Publishing Company, Westport CT.
 Jugenheimer, R.W. 1976. Corn: Improvement, Seed Production, and Uses. John Wiley & Sons, New York.
 McGee, D.C. 1988. Maize Diseases. APS Press, St. Paul, MN. 150 pp
 Munsell Color Chart for Plant Tissues. Macbeth, PO Box 230, Newburgh, N.Y. 12551-0230
 The Mutants of Maize. 1968. Crop Science Society of America. Madison, WI.
 Shurtleff, M.C. 1980. Compendium of Corn Diseases. APS Press, St. Paul, MN. 105 pp
 Sprague, G.F., and J.W. Dudley (Editors). 1988. Corn and Corn Improvement, Third Edition. Agronomy Monograph 18. ASA, CSSA, SSSA, Madison WI
 Stringfield, G.H. Maize Inbred Lines of Ohio. Ohio A.E.S., Bul. 831, 1959.
 U.S. Department of Agriculture. 1936, 1937, Yearbook.

COMMENTS (eg. State how heat units were calculated, standard inbred seed source, and/or where data was collected. Continue in Exhibit D):

Additional Description of the Inbred

Exhibit D:

JCR503 is a late season field corn inbred line that flowers similar to LH195. It is an excellent seed parent.

JCR503 has shown excellent yield potential in several hybrid combinations involving non-stiff stalk inbreds. Most hybrids involving JCR503 are adapted to the central and southern region of the corn-belt. Compared to LH195 hybrids, those crosses involving JCR503 have higher yield, and slightly more harvest moisture. LH195 crosses have less pre-anthesis brittle snapping and better roots compared to JCR503.

JCR503 INBRED DESCRIPTION

JCR503 is a corn inbred line developed from the cross of LH195 x SR108. Standard pedigree ear-row selection method was used. Selection criteria used during the development of JCR503 were: yield; high plant density tolerance; stalk and root strength; late plant greenness; and disease tolerance. Testcrosses with unrelated inbred lines were made and evaluated over multiple locations and years. Many sublines within this population had good hybrid performance across several testers. Hybrids involving this line are best adapted to the Central and Southern cornbelt.

U.S. DEPARTMENT OF AGRICULTURE
AGRICULTURAL MARKETING SERVICE

EXHIBIT E
STATEMENT OF THE BASIS OF OWNERSHIP

The following statements are made in accordance with the Privacy Act of 1974 (5 U.S.C. 552a) and the Paperwork Reduction Act (PRA) of 1995.

Application is required in order to determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421). Information is held confidential until certificate is issued (7 U.S.C. 2426).

1. NAME OF APPLICANT(S) The J. C. Robinson Seed Company	2. TEMPORARY DESIGNATION OR EXPERIMENTAL NUMBER	3. VARIETY NAME JCR503
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP Code, and Country) 100 J. C. Robinson Seed Company 100 J. C. Robinson Blvd. PO Box A Waterloo, NE 68069	5. TELEPHONE (include area code) (402)289-0252	6. FAX (include area code) (402_779-4370
	7. PVPO NUMBER 200100014	

8. Does the applicant own all rights to the variety? Mark an "X" in appropriate block. If no, please explain. YES NO

9. Is the applicant (individual or company) a U.S. national or U.S. based company? YES NO
If no, give name of country

10. Is the applicant the original owner? YES NO If no, please answer one of the following:
a. If original rights to variety were owned by individual(s), is (are) the original owner(s) a U.S. national(s)?
 YES NO If no, give name of country
b. If original rights to variety were owned by a company(ies), is(are) the original owner(s) a U.S. based company?
 YES NO If no, give name of country

11. Additional explanation of ownership (if needed, use reverse for extra space):

The variety for which Plant Variety Protection is hereby sought was developed by Calvin Treat, and David Nelson, employees of The J. C. Robinson Seed Company. By agreement between the employees and The J. C. Robinson Seed Company all rights to any invention, discovery, or development made by the employees while employed by The J. C. Robinson Seed Company are assigned to The J. C. Robinson Seed Company with no rights of any kind retained by the employees.

PLEASE NOTE:

Plant variety protection can be afforded only to owners (not licensees) who meet one of the following criteria:

1. If the rights to the variety are owned by the original breeder, that person must be a U.S. national, national of a UPOV member country, or national of a country which affords similar protection to nationals of the U.S. for the same genus and species.
2. If the rights to the variety are owned by the company which employed the original breeder(s), the company must be U.S. based, owned by nationals of a UPOV member country, or owned by nationals of a country which affords similar protection to nationals of the U.S. for the same genus and species.
3. If the applicant is an owner who is not the original owner, both the original owner and the applicant must meet one of the above criteria.

The original breeder/owner may be the individual or company who directed final breeding. See Section 41(a)(2) of the Plant Variety Protection Act for definition.

According to the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 10 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and marital or familial status. (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint, write the secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C. 20250, or call 1-800-245-6340 (voice) or (202)720-1127 (TDD). USDA is an equal employment opportunity employer.