

Cotton Varieties Planted 2014 Crop



***U. S. DEPARTMENT OF AGRICULTURE
Agricultural Marketing Service - Cotton Program
Memphis, Tennessee
September 12, 2014***

mp_cn833

Cotton Varieties Planted, United States 2014 Crop

The Deltapine brand of Upland cottonseed was the most popular planted in the United States for the 2014-2015 season, according to the USDA, Agricultural Marketing Service's Cotton and Tobacco Program. The Bayer CropScience Fibermax brand was the second most popular followed by Phytogen, Bayer CropScience Stoneville, Americot, Dyna-Gro, and All-Tex.

Transgenic varieties - genetically engineered varieties resistant to worms, herbicides, or both - accounted for about 99.50 percent of the Upland cotton planted in the United States in 2014. Usage of transgenic varieties in 2014 was reported at 100 percent in Alabama, Arkansas, Arizona, Florida, Kansas, Louisiana, Missouri, New Mexico, North Carolina, Oklahoma, Tennessee, and Virginia. Other states planted from 97.4 - 99.9 percent transgenic.

Deltapine brand varieties were the most popular planted in 2014, accounting for 29.9 percent of the United States acreage. This brand accounted for 47.2 percent of the acreage planted in the southeastern states (Alabama, Florida, Georgia, North Carolina, South Carolina, and Virginia). It accounted for about 20.2 percent in the south central states (Arkansas, Louisiana, Mississippi, Missouri, and Tennessee), 25.0 percent in the southwestern states (Texas, Oklahoma, and Kansas), and 37.1 percent in the western states (Arizona, California, and New Mexico). Deltapine's most popular varieties were DP 1044 B2RF, DP 1252 B2RF, DP 1050 B2RF, and DP 1219 B2RF, accounting respectively for 6.7, 3.9, 3.4, and 2.9 percent of the U.S. Upland cotton acreage.

Bayer CropScience Fibermax brand varieties were the second most popular planted in 2014, accounting for 22.7 percent of the United States acreage. These varieties accounted for 8.8 percent of the acreage planted in the southeastern states, 6.5 percent in the south central states, 32.0 percent in the southwestern states, and 17.3 percent in the western states. The most popular Bayer CropScience Fibermax varieties were FM 1944 GLB2, FM 2484 B2F, FM 2011 GT, and FM 2989 GLB2, accounting respectively for about 6.0, 4.4, 2.7, and 1.8 percent of the United States acreage planted to Upland cotton.

Phytogen brand varieties were the third most popular planted in 2014, accounting for 15.3 percent of the United States acreage. They accounted for 24.6 percent of the acreage planted in the southeastern states, 24.5 percent of the acreage in the south central states, 9.4 percent in the southwestern states and 20.2 percent in the western states. The most popular Phytogen brand varieties were PHY 499 WRF, PHY 375 WRF, and PHY 339 WRF, accounting respectively for 9.3, 2.1 and 1.1 percent of the United States acreage planted to Upland cotton.

Bayer CropScience Stoneville brand varieties were the fourth most popular planted in 2014. These varieties accounted for 12.4 percent of the acreage planted. They accounted for 15.6 percent of the acreage planted in the southeastern states, 43.4 percent of the acreage in the south central states, 3.8 percent in the southwestern states and 17.0 percent in the western states. The most popular Bayer CropScience Stoneville varieties were ST 4946 GLB2, ST 6448 GLB2, and ST 5289 GLT accounting respectively for 6.7, 3.0, and .6 percent of the United States acreage planted to Upland cotton.

Americot brand varieties were the fifth most popular and accounted for about 12.3 percent of the U.S. acreage planted in 2014. Dyna-Gro varieties were the sixth most popular and accounted for about 4.1 percent of the 2014 cotton acreage. All-Tex varieties were the seventh most popular and accounted for about 2.4 percent of the 2014 cotton acreage.

Phytogen was the most popular brand of American Pima varieties planted in 2014. Phytogen variety PHY 805 RF accounted for 38.0 percent of the United States Pima acreage. Phytogen's PHY 811 RF was the second most planted American Pima variety and accounted for 24.6 percent of the U.S. crop. Phytogen's PHY 802 RF was the next most popular variety and accounted for 17.9 percent of the U.S. Pima acreage.

Bayer CropScience FM 958, AFD 2485, and All-Tex 7A21 were the predominate varieties planted by organic cotton producers. Other varieties planted by organic producers include All-Tex LA122, Seed Source Genetics CT 210, and Downer Cotton Genetics DCG 1374.

Estimates of the percentage of the various varieties of cotton planted in the United States for 2014 were based on informal surveys made by the Cotton and Tobacco Program Classing Offices. Those surveyed included ginners, seed dealers, extension agents, and other knowledgeable sources.

Estimated percentage of Upland cotton planted to leading specified varieties, by growth area,

Variety	US	Southeast	South Central	Southwest	Far West
PHY 499 WRF Phytogen	9.25%	14.44%	14.46%	6.17%	4.96%
ST 4946 GLB2 Bayer CropScience - Stoneville	6.70%	2.50%	29.92%	3.05%	5.93%
DP 1044 B2RF Deltapine	6.67%	1.30%	0.42%	9.90%	15.28%
FM 1944 GLB2 Bayer CropScience - FiberMax	6.04%	7.78%	6.50%	5.46%	-
FM 2484 B2F Bayer CropScience - FiberMax	4.38%	-	-	7.19%	2.02%
NG 1511 B2RF Americot	4.11%	0.75%	2.90%	5.85%	0.12%
DP 1252 B2RF Deltapine	3.93%	14.97%	-	0.64%	0.12%
DP 1050 B2RF Deltapine	3.42%	14.40%	0.04%	-	0.04%
ST 6448 GLB2 Bayer CropScience - Stoneville	3.02%	12.08%	0.61%	0.04%	2.42%
DP 1219 B2RF Deltapine	2.95%	0.09%	*	4.70%	4.49%
FM 2011 GT Bayer CropScience - FiberMax	2.71%	-	-	4.49%	0.19%
DP 1321 B2RF Deltapine	2.69%	0.91%	3.16%	3.36%	0.55%
DG 2570 B2RF Dyna-Gro	2.58%	0.03%	1.47%	3.89%	0.97%
DP 1137 B2RF Deltapine	2.56%	9.65%	1.90%	-	0.54%
NG 4111 RF Americot	2.29%	0.05%	-	3.77%	-
DP 0912 B2RF Deltapine	2.15%	2.21%	7.80%	0.66%	6.89%
PHY 375 WRF Phytogen	2.07%	3.00%	5.80%	0.80%	3.32%
FM 2989 GLB2 Bayer CropScience - FiberMax	1.76%	-	-	2.79%	3.71%
FM 9250 GL Bayer CropScience - FiberMax	1.57%	0.03%	-	2.58%	0.12%
DP 1212 B2RF Deltapine	1.32%	-	0.12%	2.16%	-

**Estimated percentage of Upland cotton planted to leading specified varieties, by growth area,
2014 crop**

Brand	US	Southeast	South Central	Southwest	Far West
Deltapine	29.86%	47.23%	20.15%	25.01%	37.14%
Bayer CropScience - FiberMax	22.65%	8.84%	6.50%	31.98%	17.30%
Phytogen	15.31%	24.56%	24.54%	9.38%	20.23%
Bayer CropScience - Stoneville	12.38%	15.63%	43.36%	3.82%	17.04%
Americot	12.25%	2.70%	3.46%	18.46%	0.12%
Dyna-Gro	4.12%	0.10%	1.48%	6.42%	0.97%
All-Tex	2.39%	-	-	3.96%	-
Miscellaneous	0.63%	0.36%	0.35%	0.57%	7.19%
Croplan Genetics	0.39%	0.58%	0.12%	0.39%	-
Seed Source Genetics	0.01%	0.01%	0.04%	-	-

Table 1. Estimated percentage of upland cotton planted to specified varieties, by states and United States, crop of 2013

	AL	AR	AZ	CA	FL	GA	KS	LA	MO	MS	NC	NM	OK	SC	TN	TX	VA	US
All-Tex																		
1203	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.01	-	0.01
Apex B2RF	-	-	-	-	-	-	-	-	-	-	-	-	0.95	-	-	-	-	0.02
Edge B2RF	-	-	-	-	-	-	-	-	-	-	-	-	0.72	-	-	1.97	-	1.15
Epic RF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1.74	-	1.00
Nitro-44 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.31	-	0.18
Xpress	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.03	-	0.02
7A21	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.01	-	*
Excess	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.01	-	*
All-Tex Total	-	-	-	-	-	-	-	-	-	-	-	-	1.67	-	-	4.07	-	2.39
Americot																		
AM UA48	-	-	-	-	-	2.59	-	-	-	-	-	-	-	0.07	-	-	-	0.34
AM 1532 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	0.80	-	-	1.08	-	0.64
AM 1550 B2RF	-	0.02	-	-	-	0.54	-	-	0.92	0.66	0.40	-	-	0.16	1.51	0.01	0.31	0.18
NG 1511 B2RF	-	2.74	2.51	-	-	0.64	-	-	9.68	0.27	1.23	1.00	12.40	1.17	3.82	5.63	1.93	4.14
NG 1551 RF	-	-	-	-	-	-	22.79	-	-	-	0.18	-	0.45	0.13	-	0.11	0.33	0.18
NG 2549 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	6.01	-	-	0.86	-	0.62
NG 1572 RF	-	-	-	-	-	-	12.00	-	-	-	-	-	-	-	-	0.33	-	0.24
NG 2051 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2.22	-	1.28
NG 3348 B2RF	-	-	0.06	-	-	-	0.57	-	-	-	-	-	0.89	-	-	1.75	-	1.03
NG 4010 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.53	-	0.31
NG 4111 RF	0.37	-	-	-	-	-	-	-	-	-	-	-	2.89	-	-	3.83	-	2.29
NG 4012 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	0.03	-	-	1.50	-	0.87
NG 5315 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	-	0.07	-	-	-	*
NG 3306 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.28	-	0.16
Americot Total	0.37	2.75	2.57	-	-	3.78	35.36	-	10.60	0.93	1.81	1.00	23.47	1.61	5.33	18.14	2.57	12.29
Bayer CropScience - FiberMax																		
Acala Daytona RF	-	-	-	31.78	-	-	-	-	-	-	-	-	-	-	-	-	-	0.17
FM 832	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.20	-	0.11
FM 835 LLB2	-	-	-	-	-	0.05	-	-	-	-	-	-	-	-	-	0.31	-	0.19
FM 1740 B2F	3.28	-	-	-	-	0.27	-	-	-	-	0.09	4.50	9.67	-	-	1.54	-	1.26
FM 1845 LLB2	-	-	-	-	-	0.68	-	-	-	-	-	-	-	-	-	1.03	-	0.69
FM 1944 GLB2	1.63	15.24	0.63	-	-	5.43	10.14	2.09	7.09	0.17	18.27	-	4.51	9.97	8.48	5.46	12.10	6.04
FM 2011 GT	-	-	-	0.04	-	-	0.86	-	-	-	-	1.50	-	-	-	4.68	-	2.71
FM 2484 B2F	-	-	0.84	-	-	-	-	-	-	-	-	10.07	1.11	-	-	7.47	-	4.38
FM 2989 GLB2	-	-	3.90	-	-	-	-	-	-	-	-	8.86	-	-	-	2.92	-	1.76
FM 8270 GLB2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1.15	-	0.66
FM 9063 B2F	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1.67	-	0.97
FM 9170 B2F	-	-	-	-	-	-	-	-	-	-	-	-	0.95	-	-	0.36	-	0.23
FM 9180 B2F	-	-	1.15	-	-	-	-	-	-	-	-	-	0.40	-	-	2.08	-	1.23
FM 9250 GL	-	-	-	0.48	-	0.05	-	-	-	-	-	-	-	-	-	2.69	-	1.57
FM 1830 GLT	-	-	-	-	-	-	7.07	-	-	-	-	-	-	-	-	0.25	-	0.17
FM 2334 GLT	-	-	3.72	-	-	-	-	-	-	-	-	-	-	-	-	0.81	-	0.52
FM 1320 GL	-	-	-	0.24	-	-	-	-	-	-	-	-	-	-	-	-	-	*
FM 2322 GL	-	-	-	0.16	-	-	-	-	-	-	-	-	-	-	-	-	-	*
Bayer CropScience - FiberMax Total	4.91	15.24	10.24	32.70	-	6.48	18.07	2.09	7.09	0.17	18.36	24.93	16.65	9.97	8.48	32.63	12.10	22.66

Table 1. Estimated percentage of upland cotton planted to specified varieties, by states and United States, crop of 2013

	AL	AR	AZ	CA	FL	GA	KS	LA	MO	MS	NC	NM	OK	SC	TN	TX	VA	US
Bayer CropScience - Stoneville																		
ST 4145 LLB2	-	1.27	-	-	-	-	-	-	-	-	0.53	-	-	0.93	0.84	-	0.33	0.11
ST 4288 B2F	0.19	-	-	-	-	-	-	-	-	0.05	-	-	0.95	-	-	-	-	0.03
ST 4498 B2RF	1.40	-	7.98	2.35	-	0.37	-	-	-	-	-	-	-	-	-	-	-	0.21
ST 4946 GLB2	0.34	48.14	7.10	5.37	-	0.33	-	2.10	45.88	27.00	10.32	2.64	-	2.30	24.51	3.18	7.64	6.70
ST 5288 B2F	-	0.85	-	-	-	-	-	21.52	0.71	0.89	0.51	-	-	-	-	-	-	0.41
ST 5458 B2RF	-	0.26	0.28	-	-	-	-	-	0.05	1.30	-	-	-	-	-	0.50	-	0.35
ST 5445 LLB2	-	0.17	-	-	-	-	-	-	0.13	-	0.01	-	-	-	-	-	0.25	0.01
ST 5458 B2F	-	8.42	1.14	-	-	-	-	-	4.71	0.25	-	-	-	-	0.76	0.14	-	0.50
ST 6448 GLB2	12.98	0.58	3.18	0.97	-	14.61	-	2.67	-	0.75	3.88	2.14	-	14.76	-	0.04	0.97	3.02
ST 4747 GLB2	0.07	2.60	0.35	0.97	0.51	-	-	-	0.97	1.15	1.92	9.00	-	0.54	0.08	0.09	0.74	0.34
ST 5032 GLT	-	1.74	-	-	-	0.05	-	-	1.85	-	-	-	-	-	-	-	-	0.10
ST 5289 GLT	-	1.17	0.51	2.35	-	-	-	4.72	0.25	12.55	0.28	-	-	0.03	-	-	0.59	0.60
Bayer CropScience - Stoneville Total	14.97	65.20	20.53	12.00	0.51	15.36	-	31.00	54.55	43.95	17.45	13.79	0.95	18.57	26.19	3.95	10.52	12.38
Concho																		
257	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.01	-	0.01
Concho Total	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.01	-	0.01
Croplan Genetics																		
CG 3220 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.76	0.13	-	0.09
CG 3787 B2RF	-	-	-	-	1.69	0.92	-	-	-	-	-	-	-	-	-	0.28	-	0.30
Croplan Genetics Total	-	-	-	-	1.69	0.92	-	-	-	-	-	-	-	-	0.76	0.41	-	0.39
Deltapine																		
DP 0912 B2RF	2.44	6.17	10.48	2.35	0.56	1.71	23.64	4.15	14.48	5.62	4.16	-	14.30	2.36	11.96	-	1.07	2.15
DP 1028 B2RF	3.28	-	-	-	1.01	0.57	-	-	-	-	5.51	-	-	0.41	-	-	3.85	0.47
DP 1034 B2RF	0.23	-	-	-	-	0.23	-	-	-	4.22	1.50	-	2.84	0.21	1.51	-	0.13	0.35
DP 1044 B2RF	2.41	-	21.29	3.52	-	1.73	-	0.05	0.40	1.35	-	14.50	13.63	-	-	9.81	-	6.67
DP 1050 B2RF	10.36	-	-	0.17	58.54	18.48	-	0.09	-	0.11	0.38	-	-	1.61	-	-	-	3.42
DP 1133 B2RF	5.02	2.11	0.19	-	-	0.17	-	19.80	-	0.78	0.67	-	-	0.18	-	0.04	0.64	0.64
DP 1137 B2RF	10.28	-	0.78	0.28	20.22	10.95	-	0.19	-	2.84	6.68	-	-	7.33	-	-	0.97	2.56
DP 1212 B2RF	0.42	0.02	-	-	-	-	1.43	-	-	-	-	-	0.18	-	0.11	2.24	-	1.32
DP 1219 B2RF	0.23	0.02	7.45	-	-	0.10	-	-	-	-	-	-	4.47	-	-	4.73	-	2.95
DP 1252 B2RF	4.93	-	-	0.47	9.66	23.63	-	-	-	-	0.58	-	-	7.61	-	0.67	-	3.93
DP 1311 B2RF	-	1.76	0.74	-	-	-	-	1.05	1.92	0.23	0.13	-	-	-	4.58	0.16	-	0.33
DP 1321 B2RF	6.21	0.34	0.92	-	3.37	0.53	14.79	1.74	0.19	4.62	1.64	-	-	1.03	1.34	3.40	1.36	2.69
DP 1359 B2RF	0.12	-	13.29	0.71	-	-	-	-	-	-	-	-	1.80	-	-	1.68	-	1.19
DP 1454NR B2RF	0.40	-	-	-	2.25	0.88	-	-	-	-	0.02	-	-	0.51	-	1.57	-	1.07
DP 1410 B2RF	-	0.03	0.04	-	-	-	0.36	-	-	-	-	-	-	-	-	0.12	-	0.07
DP 1441 RF	-	-	0.06	2.51	-	-	-	-	-	0.96	-	-	-	-	-	-	-	0.05
Deltapine Total	46.33	10.45	55.24	10.01	95.62	58.99	40.21	27.07	17.00	20.75	21.29	14.50	37.21	21.24	19.50	24.42	8.03	29.87
Dyna-Gro																		
DG 2285 B2RF	0.44	-	-	-	-	-	-	0.07	-	-	-	-	-	-	-	1.32	-	0.78
DG 2450 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	0.72	-	-	1.29	-	0.76
DG 2570 B2RF	0.25	0.63	-	-	-	-	-	*	5.89	0.48	-	7.86	12.17	-	1.51	3.60	-	2.58
DG 2610 B2RF	-	-	-	-	-	-	-	*	-	-	-	-	-	-	-	-	-	*
Dyna-Gro Total	0.68	0.63	-	-	-	-	-	0.08	5.89	0.48	-	7.86	12.89	-	1.51	6.21	-	4.12

Table 1. Estimated percentage of upland cotton planted to specified varieties, by states and United States, crop of 2013

	AL	AR	AZ	CA	FL	GA	KS	LA	MO	MS	NC	NM	OK	SC	TN	TX	VA	US
Phytogen																		
PHY 72	-	-	-	0.42	-	-	-	-	-	-	-	-	-	-	-	-	-	*
PHY 339 WRF	1.08	0.28	0.56	-	1.69	0.06	2.21	4.08	-	4.67	7.05	7.93	-	3.63	0.25	0.43	12.13	1.07
PHY 367 WRF	0.07	-	0.85	1.05	-	-	-	-	-	-	0.42	1.07	-	1.47	-	0.43	2.22	0.34
PHY 375 WRF	8.12	0.02	0.69	3.52	-	0.60	4.14	1.45	2.92	1.16	7.18	8.93	2.39	2.22	23.78	0.76	18.43	2.07
PHY 440 W	0.01	-	-	-	-	0.33	-	-	-	0.51	-	-	-	-	-	*	-	0.06
PHY 499 WRF	15.32	5.11	2.73	5.10	0.51	8.49	-	33.67	1.88	21.76	22.54	14.43	0.48	36.76	11.73	6.42	30.17	9.25
PHY 575 WRF	4.39	-	-	-	-	0.80	-	-	-	-	0.02	-	-	0.77	-	0.01	-	0.28
PHY 725 RF	-	-	1.39	27.56	-	-	-	-	-	-	-	-	-	-	-	-	-	0.17
PHY 755 WRF	-	-	-	4.59	-	-	-	-	-	1.78	-	-	-	-	-	-	-	0.09
PHY 333 WRF	2.84	0.21	0.38	-	-	0.40	-	0.57	0.04	1.70	2.55	1.07	-	1.19	2.48	-	3.44	0.45
PHY 417 WRF	-	0.02	-	0.08	-	-	-	-	-	-	0.20	-	-	0.06	-	1.56	-	0.91
PHY 427 WRF	-	-	-	0.07	-	3.40	-	-	-	0.71	0.51	-	-	0.29	-	-	0.33	0.50
PHY 495 W3RF	-	0.10	-	-	-	-	-	-	-	-	0.02	-	-	0.03	-	-	0.02	*
PHY 599 WRF	0.82	-	-	-	-	0.34	-	-	-	-	-	-	2.23	-	-	-	-	0.12
PHY 222 WRF	-	-	0.06	-	-	-	-	-	-	-	-	-	-	-	-	-	-	*
Phytogen Total	32.66	5.73	6.66	42.39	2.19	14.42	6.36	39.77	4.84	32.27	40.49	33.43	5.09	46.42	38.23	9.61	66.75	15.31
Seed Source Genetics																		
CT 110	-	-	-	-	-	-	-	-	-	0.15	-	-	-	-	-	-	-	0.01
CT 210	-	-	-	-	-	0.02	-	-	-	-	-	-	-	-	-	-	-	*
Seed Source Genetics Total	-	-	-	-	-	0.02	-	-	-	0.15	-	-	-	-	-	-	-	0.01
Miscellaneous																		
Misc.	0.07	-	4.75	2.90	-	0.04	-	-	0.04	1.30	0.60	4.50	2.07	2.18	-	0.55	0.03	0.59
Miscellaneous Total	0.07	-	4.75	2.90	-	0.04	-	-	0.04	1.30	0.60	4.50	2.07	2.18	-	0.55	0.03	0.59
Transgenic	100.00%	100.00%	100.00%	99.58%	100.00%	97.39%	100.00%	100.00%	100.00%	99.85%	100.00%	100.00%	100.00%	99.93%	100.00%	99.74%	100.00%	99.50%
Conventional	0.00%	0.00%	0.00%	0.42%	0.00%	2.61%	0.00%	0.00%	0.00%	0.15%	0.00%	0.00%	0.00%	0.07%	0.00%	0.25%	0.00%	0.49%

Table 2. Estimated percentage of upland cotton planted to specified varieties, by Classing Office and United States, crop of 2013

Variety	US	Abilene				Corpus	Dumas			Florence			
	Total	KS	OK	TX	Classing Office	TX	AR	MS	Classing Office	NC	SC	VA	Classing Office
PHY 499 WRF Phytogen	9.25%	-	0.48%	0.69%	0.60%	21.06%	11.32%	20.58%	18.68%	22.54%	36.76%	30.17%	27.94%
ST 4946 GLB2 Bayer CropScience - Stoneville	6.70%	-	-	1.98%	1.37%	0.90%	43.29%	29.97%	32.70%	10.32%	2.30%	7.64%	7.45%
DP 1044 B2RF Deltapine	6.67%	-	13.63%	21.55%	18.50%	10.41%	-	-	-	-	-	-	-
FM 1944 GLB2 Bayer CropScience - FiberMax	6.04%	10.14%	4.51%	12.98%	10.66%	16.73%	2.57%	0.20%	0.69%	18.27%	9.97%	12.10%	14.94%
FM 2484 B2F Bayer CropScience - FiberMax	4.38%	-	1.11%	0.90%	0.91%	-	-	-	-	-	-	-	-
NG 1511 B2RF Americot	4.14%	-	12.40%	5.25%	6.86%	0.03%	2.31%	0.31%	0.73%	1.23%	1.17%	1.93%	1.28%
DP 1252 B2RF Deltapine	3.93%	-	-	-	-	3.21%	-	-	-	0.58%	7.61%	-	2.78%
DP 1050 B2RF Deltapine	3.42%	-	-	-	-	-	-	-	-	0.38%	1.61%	-	0.74%
ST 6448 GLB2 Bayer CropScience - Stoneville	3.02%	-	-	-	-	0.18%	-	0.87%	0.69%	3.88%	14.76%	0.97%	7.07%
DP 1219 B2RF Deltapine	2.95%	-	4.47%	9.69%	7.89%	9.99%	-	-	-	-	-	-	-
FM 2011 GT Bayer CropScience - FiberMax	2.71%	0.86%	-	1.45%	1.05%	-	-	-	-	-	-	-	-
DP 1321 B2RF Deltapine	2.69%	14.79%	-	3.45%	3.09%	-	0.41%	3.52%	2.88%	1.64%	1.03%	1.36%	1.41%
DG 2570 B2RF Dyna-Gro	2.58%	-	12.17%	7.42%	8.30%	0.39%	-	-	-	-	-	-	-
DP 1137 B2RF Deltapine	2.56%	-	-	-	-	-	-	3.27%	2.59%	6.68%	7.33%	0.97%	6.28%
NG 4111 RF Americot	2.29%	-	2.89%	1.17%	1.56%	-	-	-	-	-	-	-	-
DP 0912 B2RF Deltapine	2.15%	23.64%	14.30%	-	4.79%	-	24.38%	6.06%	9.82%	4.16%	2.36%	1.07%	3.25%
PHY 375 WRF Phytogen	2.07%	4.14%	2.39%	-	0.81%	3.00%	-	1.33%	1.06%	7.18%	2.22%	18.43%	6.78%
FM 2989 GLB2 Bayer CropScience - FiberMax	1.76%	-	-	4.05%	2.82%	2.52%	-	-	-	-	-	-	-
FM 9250 GL Bayer CropScience - FiberMax	1.57%	-	-	0.50%	0.35%	-	-	-	-	-	-	-	-
DP 1212 B2RF Deltapine	1.32%	1.43%	0.18%	-	0.11%	2.31%	-	-	-	-	-	-	-
NG 2051 B2RF Americot	1.28%	-	-	-	-	-	-	-	-	-	-	-	-
FM 1740 B2F Bayer CropScience - FiberMax	1.26%	-	9.67%	4.80%	5.84%	1.63%	-	-	-	0.09%	-	-	0.05%
FM 9180 B2F Bayer CropScience - FiberMax	1.23%	-	0.40%	6.94%	4.93%	-	-	-	-	-	-	-	-

Table 2. Estimated percentage of upland cotton planted to specified varieties, by Classing Office and United States, crop of 2013

Variety	US	Lamesa			Lubbock	Macon				Memphis					
	Total	NM /1	TX	Classing Office	TX	AL	FL	GA	Classing Office	AL	AR	MO	MS	TN	Classing Office
PHY 499 WRF Phytogen	9.25%		1.55%	1.59%	3.60%	10.93%	0.51%	8.49%	8.32%	24.21%	3.05%	1.88%	37.47%	11.73%	9.34%
ST 4946 GLB2 Bayer CropScience - Stoneville	6.70%		5.11%	5.11%	3.40%	-	-	0.33%	0.26%	1.02%	49.76%	45.88%	8.91%	24.51%	33.75%
DP 1044 B2RF Deltapine	6.67%		10.42%	10.42%	6.45%	3.60%	-	1.73%	1.88%	-	-	0.40%	13.49%	-	0.68%
FM 1944 GLB2 Bayer CropScience - FiberMax	6.04%		1.62%	1.61%	0.58%	1.43%	-	5.43%	4.53%	2.02%	19.46%	7.09%	-	8.48%	10.07%
FM 2484 B2F Bayer CropScience - FiberMax	4.38%		8.49%	8.48%	11.99%	-	-	-	-	-	-	-	-	-	-
NG 1511 B2RF Americot	4.14%		7.70%	7.71%	7.20%	-	-	0.64%	0.51%	-	2.88%	9.68%	-	3.82%	4.45%
DP 1252 B2RF Deltapine	3.93%		-	-	-	7.35%	9.66%	23.63%	20.49%	-	-	-	-	-	-
DP 1050 B2RF Deltapine	3.42%		-	-	-	15.46%	58.54%	18.48%	20.60%	-	-	-	-	-	-
ST 6448 GLB2 Bayer CropScience - Stoneville	3.02%		-	-	-	19.37%	-	14.61%	14.35%	-	0.77%	-	-	-	0.22%
DP 1219 B2RF Deltapine	2.95%		2.76%	2.76%	2.23%	0.34%	-	0.10%	0.13%	-	0.02%	-	-	-	0.01%
FM 2011 GT Bayer CropScience - FiberMax	2.71%		3.41%	3.43%	8.47%	-	-	-	-	-	-	-	-	-	-
DP 1321 B2RF Deltapine	2.69%		4.67%	4.66%	4.28%	0.34%	3.37%	0.53%	0.68%	18.15%	0.32%	0.19%	13.49%	1.34%	3.48%
DG 2570 B2RF Dyna-Gro	2.58%		5.83%	5.84%	2.93%	0.37%	-	-	0.05%	-	0.84%	5.89%	-	1.51%	2.22%
DP 1137 B2RF Deltapine	2.56%		-	-	-	8.34%	20.22%	10.95%	11.18%	14.23%	-	-	-	-	1.89%
NG 4111 RF Americot	2.29%		5.27%	5.26%	5.44%	0.56%	-	-	0.08%	-	-	-	-	-	-
DP 0912 B2RF Deltapine	2.15%		-	-	-	2.45%	0.56%	1.71%	1.74%	2.41%	0.10%	14.48%	3.54%	11.96%	7.58%
PHY 375 WRF Phytogen	2.07%		0.14%	0.16%	0.13%	5.86%	-	0.60%	1.29%	12.73%	0.02%	2.92%	-	23.78%	8.84%
FM 2989 GLB2 Bayer CropScience - FiberMax	1.76%		2.36%	2.39%	3.17%	-	-	-	-	-	-	-	-	-	-
FM 9250 GL Bayer CropScience - FiberMax	1.57%		2.91%	2.91%	4.38%	-	-	0.05%	0.04%	-	-	-	-	-	-
DP 1212 B2RF Deltapine	1.32%		1.11%	1.11%	3.42%	-	-	-	-	1.28%	0.02%	-	-	0.11%	0.21%
NG 2051 B2RF Americot	1.28%		0.73%	0.73%	4.72%	-	-	-	-	-	-	-	-	-	-
FM 1740 B2F Bayer CropScience - FiberMax	1.26%		1.60%	1.60%	0.71%	4.90%	-	0.27%	0.90%	-	-	-	-	-	-
FM 9180 B2F Bayer CropScience - FiberMax	1.23%		0.19%	0.19%	3.11%	-	-	-	-	-	-	-	-	-	-

Table 2. Estimated percentage of upland cotton planted to specified varieties, by Classing Office and United States, crop of 2013

Variety	US	Rayville			Visalia				Classing Office
	Total	LA	MS /1	Classing Office	AZ	CA	NM	TX /1	
PHY 499 WRF Phytogen	9.25%	33.67%		31.55%	2.73%	5.10%	13.81%		4.77%
ST 4946 GLB2 Bayer CropScience - Stoneville	6.70%	2.10%		2.04%	7.10%	5.37%	2.38%		5.94%
DP 1044 B2RF Deltapine	6.67%	0.05%		0.04%	21.29%	3.52%	15.00%		15.55%
FM 1944 GLB2 Bayer CropScience - FiberMax	6.04%	2.09%		1.94%	0.63%	-	-		0.38%
FM 2484 B2F Bayer CropScience - FiberMax	4.38%	-		-	0.84%	-	11.19%		2.04%
NG 1511 B2RF Americot	4.14%	-		-	2.51%	-	-		1.53%
DP 1252 B2RF Deltapine	3.93%	-		-	-	0.47%	-		0.12%
DP 1050 B2RF Deltapine	3.42%	0.09%		0.34%	-	0.17%	-		0.04%
ST 6448 GLB2 Bayer CropScience - Stoneville	3.02%	2.67%		2.48%	3.18%	0.97%	2.38%		2.45%
DP 1219 B2RF Deltapine	2.95%	-		-	7.45%	-	-		4.55%
FM 2011 GT Bayer CropScience - FiberMax	2.71%	-		-	-	0.04%	-		0.01%
DP 1321 B2RF Deltapine	2.69%	1.74%		2.10%	0.92%	-	-		0.56%
DG 2570 B2RF Dyna-Gro	2.58%	*		1.11%	-	-	7.62%		0.86%
DP 1137 B2RF Deltapine	2.56%	0.19%		0.18%	0.78%	0.28%	-		0.55%
NG 4111 RF Americot	2.29%	-		-	-	-	-		-
DP 0912 B2RF Deltapine	2.15%	4.15%		3.86%	10.48%	2.35%	-		6.98%
PHY 375 WRF Phytogen	2.07%	1.45%		1.35%	0.69%	3.52%	8.81%		3.24%
FM 2989 GLB2 Bayer CropScience - FiberMax	1.76%	-		-	3.90%	-	7.62%		3.51%
FM 9250 GL Bayer CropScience - FiberMax	1.57%	-		-	-	0.48%	-		0.12%
DP 1212 B2RF Deltapine	1.32%	-		-	-	-	-		-
NG 2051 B2RF Americot	1.28%	-		-	-	-	-		-
FM 1740 B2F Bayer CropScience - FiberMax	1.26%	-		-	-	-	5.00%		0.56%
FM 9180 B2F Bayer CropScience - FiberMax	1.23%	-		-	1.15%	-	-		0.70%

Table 3. Estimated percentage of American Pima cotton acreage planted to specified varieties, by states, crop of 2014

		AZ	CA	NM \1	TX \1	US
Phytogen	PHY 805 RF	48.77%	39.60%			37.95%
Phytogen	PHY 811 RF	25.11%	25.92%			24.57%
Phytogen	PHY 802 RF	0.00%	18.49%			17.90%
Deltapine	DP 340	8.72%	4.02%			6.11%
Hazera	HA 211	0.00%	4.44%			3.71%
Miscellaneous	MISC	4.26%	3.25%			3.07%
Deltapine	DP 357	0.00%	0.43%			2.38%
Phytogen	PHY 830	3.65%	2.24%			2.18%
Deltapine	DP 358 RF	9.50%	0.28%			1.03%
Hazera	HA 1432	0.00%	1.00%			0.83%
Hazera	HA 195	0.00%	0.20%			0.17%
Hazera	HA 222	0.00%	0.12%			0.10%

/1 Individual state data withheld