

Cotton Varieties Planted 2015 Crop



***U. S. DEPARTMENT OF AGRICULTURE
Agricultural Marketing Service - Cotton and Tobacco Program
Memphis, Tennessee
September 15, 2015***

mp_cn833

Cotton Varieties Planted, United States 2015 Crop

The Deltapine brand of Upland cottonseed was the most popular planted in the United States for the 2015-2016 season, according to the USDA, Agricultural Marketing Service's Cotton and Tobacco Program. The Bayer CropScience Fibermax brand was the second most popular followed by Bayer CropScience Stoneville, Phytogen, Americot, Dyna-Gro, and All-Tex.

Transgenic varieties - genetically engineered varieties resistant to worms, herbicides, or both - accounted for about 98.8 percent of the Upland cotton planted in the United States in 2015. Usage of transgenic varieties in 2015 was reported at 100 percent in Alabama, Arkansas, Arizona, Florida, Missouri, Mississippi, New Mexico, and Tennessee. Other states planted from 95.9 - 99.6 percent transgenic.

Deltapine brand varieties were the most popular planted in 2015, accounting for 31.2 percent of the United States acreage. This brand accounted for 50.1 percent of the acreage planted in the southeastern states (Alabama, Florida, Georgia, North Carolina, South Carolina, and Virginia). It accounted for about 29.8 percent in the south central states (Arkansas, Louisiana, Mississippi, Missouri, and Tennessee), 23.7 percent in the southwestern states (Texas, Oklahoma, and Kansas), and 24.9 percent in the western states (Arizona, California, and New Mexico). Deltapine's most popular varieties were DP 1044 B2RF, DP 1219 B2RF, DP 1252 B2RF, and DP 1553 B2XF, accounting respectively for 5.6, 3.7, 3.4, and 2.0 percent of the U.S. Upland cotton acreage.

Bayer CropScience Fibermax brand varieties were the second most popular planted in 2015, accounting for 21.6 percent of the United States acreage. These varieties accounted for 3.7 percent of the acreage planted in the southeastern states, 1.0 percent in the south central states, 33.4 percent in the southwestern states, and 28.3 percent in the western states. The most popular Bayer CropScience Fibermax varieties were FM 2011 GT, FM 2484 B2F, FM 9250 GL, and FM 1944 GLB2, accounting respectively for about 4.7, 4.3, 3.8, and 2.9 percent of the United States acreage planted to Upland cotton.

Bayer CropScience Stoneville brand varieties were the third most popular planted in 2015, accounting for 16.9 percent of the United States acreage. They accounted for 13.4 percent of the acreage planted in the southeastern states, 44.7 percent of the acreage in the south central states, 12.5 percent in the southwestern states, and 17.1 percent in the western states. The most popular Bayer CropScience Stoneville brand varieties were ST 4946 GLB2, ST 4747 GLB2, and ST 6448 GLB2, accounting respectively for 10.9, 4.2, and 1.1 percent of the United States acreage planted to Upland cotton.

Phytogen brand varieties were the fourth most popular planted in 2015. These varieties accounted for 15.3 percent of the acreage planted. They accounted for 28.2 percent of the acreage planted in the southeastern states, 17.5 percent of the acreage in the south central states, 9.0 percent in the southwestern states, and 23.2 percent in the western states. The most popular Bayer Phytogen varieties were PHY 499 WRF, PHY333 WRF, and PHY 339 WRF accounting respectively for 6.1, 5.0, and 1.1 percent of the United States acreage planted to Upland cotton.

Americot brand varieties were the fifth most popular and accounted for about 6.4 percent of the U.S. acreage planted in 2015. Dyna-Gro varieties were the sixth most popular and accounted for about 4.9 percent of the 2015 cotton acreage. All-Tex varieties were the seventh most popular and accounted for about 2.3 percent of the 2015 cotton acreage.

Phytogen was the most popular brand of American Pima varieties planted in 2015. Phytogen variety PHY 805 RF accounted for 40.6 percent of the United States Pima acreage. Phytogen's PHY 811 RF was the second most planted American Pima variety and accounted for 32.8 percent of the U.S. crop. Deltapine's DP 358 RF was the next most popular variety and accounted for 9.7 percent of the U.S. Pima acreage.

AFD 2485, Bayer CropScience FM 958, All-Tex 7A21, and Deltapine DP 340 were the predominate varieties planted by organic cotton producers. Other varieties planted by organic producers include All-Tex LA122 and A102, Seed Source Genetics CT 210, and Downer Cotton Genetics DCG 1374.

Estimates of the percentage of the various varieties of cotton planted in the United States for 2015 were based on informal surveys made by the Cotton and Tobacco Program Classing Offices. Those surveyed included ginners, seed dealers, extension agents, and other knowledgeable sources.

**Estimated percentage of Upland cotton planted to leading specified varieties, by growth area,
2015 Crop**

Variety	US	Southeast	South Central	Southwest	Far West
ST 4946 GLB2 Bayer CropScience - Stoneville	10.92%	7.70%	32.11%	7.82%	7.81%
PHY 499 WRF Phytogen	6.07%	13.24%	6.37%	3.01%	5.36%
DP 1044 B2RF Deltapine	5.62%	0.72%	0.01%	8.65%	12.94%
PHY 333 WRF Phytogen	5.02%	7.78%	9.41%	3.06%	0.75%
FM 2011 GT Bayer CropScience - FiberMax	4.72%	-	-	7.86%	1.02%
FM 2484 B2F Bayer CropScience - FiberMax	4.30%	-	-	7.02%	4.83%
ST 4747 GLB2 Bayer CropScience - Stoneville	4.18%	1.42%	8.62%	4.52%	-
FM 9250 GL Bayer CropScience - FiberMax	3.82%	-	-	6.39%	-
DP 1219 B2RF Deltapine	3.70%	-	-	6.13%	1.84%
DP 1252 B2RF Deltapine	3.35%	11.81%	0.73%	0.46%	-
NG 1511 B2RF Americot	3.22%	0.50%	2.00%	4.69%	1.27%
FM 1944 GLB2 Bayer CropScience - FiberMax	2.92%	3.65%	1.02%	3.10%	0.31%
FM 2334 GLT Bayer CropScience - FiberMax	2.82%	-	-	4.53%	5.61%
DG 2570 B2RF Dyna-Gro	2.79%	0.36%	0.67%	4.30%	2.05%
DP 1553 B2XF Deltapine	1.97%	7.12%	0.01%	0.25%	1.08%
DP 1050 B2RF Deltapine	1.92%	7.59%	0.03%	-	-
DP 1538 B2XF Deltapine	1.88%	6.88%	1.09%	-	-
DP 1522 B2XF Deltapine	1.60%	1.43%	4.57%	1.06%	0.78%
DP 1137 B2RF Deltapine	1.59%	5.21%	2.14%	-	-
DP 912 B2RF Deltapine	1.57%	1.62%	3.05%	1.18%	3.01%

**Estimated percentage of Upland cotton planted to leading specified varieties, by growth area,
2015 crop**

Brand	US	Southeast	South Central	Southwest	Far West
Deltapine	31.15%	50.10%	29.81%	23.65%	24.88%
Bayer CropScience - FiberMax	21.61%	3.65%	1.02%	33.41%	28.26%
Bayer CropScience - Stoneville	16.93%	13.37%	44.66%	12.45%	17.07%
Phytogen	15.26%	28.22%	17.49%	9.04%	23.17%
Americot	6.41%	2.63%	5.26%	8.41%	1.87%
Dyna-Gro	4.85%	1.02%	1.72%	7.21%	2.94%
All-Tex	2.29%	-	0.01%	3.83%	-
Miscellaneous	1.05%	0.46%	0.03%	1.50%	1.82%
Croplan Genetics	0.43%	0.49%	-	0.50%	-
Seed Source Genetics	0.02%	0.05%	-	*	-

Table 1. Estimated percentage of upland cotton planted to specified varieties, by states and United States, crop of 2015

	AL	AR	AZ	CA	FL	GA	KS	LA	MO	MS	NC	NM	OK	SC	TN	TX	VA	US
All-Tex																		
1203	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.01	-	*
Apex B2RF	-	-	-	-	-	-	-	-	-	-	-	-	0.96	-	-	-	-	0.02
Edge B2RF	-	-	-	-	-	-	1.67	-	-	-	-	-	-	-	-	1.23	-	0.70
Epic RF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2.68	-	1.53
Nitro-44 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.04	-	0.02
LA122	-	-	-	-	-	-	-	0.13	-	-	-	-	-	-	-	-	-	*
All-Tex Total	-	-	-	-	-	-	1.67	0.13	-	-	-	-	0.96	-	-	3.95	-	2.29
Americot																		
AM UA48	-	-	-	-	-	-	-	-	-	-	0.03	-	-	0.02	-	-	-	*
AM 1532 B2RF	-	-	-	-	-	0.02	-	-	-	-	-	-	0.80	-	-	0.02	-	0.03
AM 1550 B2RF	-	-	-	-	-	0.51	-	-	0.09	-	0.01	-	-	0.02	-	-	0.27	0.07
NG 1511 B2RF	-	2.88	0.48	-	-	0.54	1.84	0.25	1.12	3.66	0.75	5.00	5.52	-	1.04	4.66	2.07	3.22
NG 1551 RF	-	-	-	-	-	-	1.28	-	0.06	-	-	-	0.16	-	-	*	-	0.01
NG 1572 RF	-	-	-	-	-	-	4.25	-	-	-	-	-	-	-	-	-	-	0.01
NG 3348 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	2.00	-	-	0.01	-	0.06
NG 4010 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	3.20	-	-	1.09	-	0.70
NG 4111 RF	0.29	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1.24	-	0.72
NG 4012 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.01	-	0.01
NG 5315 B2RF	-	-	-	-	-	0.09	-	-	-	-	-	-	-	-	-	-	-	0.01
NG 3306 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	1.60	-	-	1.16	-	0.70
NG 3405 B2XF	-	1.62	-	-	-	0.03	-	-	1.03	0.45	0.07	-	-	0.67	2.78	-	0.05	0.16
NG 3406 B2XF	-	1.96	0.97	-	-	0.01	-	-	6.62	-	1.74	-	-	7.68	4.90	-	1.89	0.61
NG 5007 B2XF	-	-	0.24	-	-	0.22	-	-	-	-	-	-	-	2.40	-	-	-	0.10
Americot Total	0.29	6.45	1.70	-	-	1.41	7.37	0.25	8.91	4.11	2.60	5.00	13.28	10.79	8.73	8.19	4.29	6.41
Bayer CropScience - FiberMax																		
FM 832	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.25	-	0.15
FM 1944 GLB2	2.20	1.39	0.63	-	1.76	2.51	9.08	-	2.52	0.15	7.81	-	2.32	2.79	1.93	3.11	5.03	2.92
FM 2011 GT	-	-	-	-	-	-	-	-	-	-	-	5.00	-	-	-	8.22	-	4.72
FM 2484 B2F	-	-	1.80	4.19	-	-	-	-	-	-	-	12.50	-	-	-	7.35	-	4.30
FM 2989 GLB2	-	-	-	-	-	-	-	-	-	-	-	10.00	-	-	-	0.79	-	0.49
FM 8270 GLB2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.08	-	0.04
FM 9180 B2F	-	-	-	-	-	-	-	-	-	-	-	-	0.48	-	-	0.02	-	0.02
FM 9250 GL	-	-	-	-	-	-	0.66	-	-	-	-	-	-	-	-	6.68	-	3.82
FM 1830 GLT	-	-	13.41	3.55	-	-	6.14	-	-	-	-	-	0.32	-	-	1.35	-	0.95
FM 2334 GLT	-	-	6.34	3.22	-	-	-	-	-	-	-	7.67	-	-	-	4.74	-	2.82
FM 1320 GL	-	-	3.97	1.31	-	-	-	-	-	-	-	-	-	-	-	-	-	0.05
FM 2322 GL	-	-	0.16	3.21	-	-	-	-	-	-	-	-	-	-	-	0.17	-	0.12
FM 1900 GLT	-	-	3.06	3.91	-	-	-	-	-	-	-	-	5.76	-	-	0.62	-	0.55
FM 2007 GLT	-	-	1.05	1.32	-	-	-	-	-	-	-	-	-	-	-	1.12	-	0.66
Bayer CropScience - FiberMax Total	2.20	1.39	30.42	20.71	1.76	2.51	15.87	-	2.52	0.15	7.81	35.17	8.88	2.79	1.93	34.52	5.03	21.61

Table 1. Estimated percentage of upland cotton planted to specified varieties, by states and United States, crop of 2015

	AL	AR	AZ	CA	FL	GA	KS	LA	MO	MS	NC	NM	OK	SC	TN	TX	VA	US
Bayer CropScience - Stoneville																		
ST 4946 GLB2	3.59	52.63	14.91	1.42	-	1.76	-	1.72	48.41	29.01	25.33	-	0.08	8.37	30.97	8.18	24.07	10.92
ST 6448 GLB2	2.20	0.35	0.40	3.30	-	3.97	-	4.97	-	0.54	2.97	-	-	7.21	-	0.12	0.33	1.14
ST 4747 GLB2	4.80	17.39	-	-	-	0.95	26.02	-	14.62	0.43	3.33	-	2.08	1.50	8.63	4.55	3.73	4.18
ST 5032 GLT	-	0.45	0.32	5.25	-	-	-	-	2.83	0.15	-	-	-	0.07	2.78	-	-	0.17
ST 5289 GLT	-	0.85	0.40	-	-	0.08	-	3.97	0.89	0.78	-	-	-	-	-	-	-	0.14
ST 5115 GLT	-	0.35	6.59	3.56	-	0.01	-	-	0.37	3.34	0.28	-	-	0.52	-	-	0.10	0.26
ST 6182 GLT	-	0.35	4.20	-	-	0.24	-	-	-	0.17	0.23	-	-	1.10	-	-	0.03	0.13
Bayer CropScience - Stoneville Total	10.59	72.37	26.83	13.54	-	7.03	26.02	10.66	67.13	34.42	32.14	-	2.16	18.78	42.38	12.85	28.26	16.93
Croplan Genetics																		
CG 3156 B2RF	-	-	-	-	-	0.21	-	-	-	-	-	-	-	-	-	-	-	0.03
CG 3787 B2RF	-	-	-	-	-	0.72	-	-	-	-	-	-	-	-	-	0.29	-	0.26
CG 3475 B2XF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.09	-	0.05
CG 3885 B2XF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.14	-	0.08
Croplan Genetics Total	-	-	-	-	-	0.93	-	-	-	-	-	-	-	-	-	0.53	-	0.43
Deltapine																		
DP 1028 B2RF	-	0.06	-	-	-	0.31	-	0.53	-	-	1.41	-	-	-	-	-	2.16	0.14
DP 1034 B2RF	0.55	-	-	-	-	0.14	-	-	-	4.53	0.49	-	-	0.16	-	-	0.03	0.24
DP 1044 B2RF	2.03	-	23.39	4.72	10.55	-	-	0.06	-	-	-	-	17.12	-	-	8.31	-	5.62
DP 1050 B2RF	8.99	-	-	-	0.84	11.70	-	0.27	-	-	-	-	-	0.41	-	-	-	1.92
DP 1133 B2RF	2.86	2.38	-	1.07	-	0.05	-	21.22	-	8.60	0.34	-	-	0.07	0.35	-	-	0.81
DP 1137 B2RF	4.39	-	-	-	17.06	7.29	-	4.35	-	2.72	1.58	-	-	1.71	-	-	0.10	1.59
DP 1212 B2RF	-	-	-	-	-	0.21	-	-	-	-	-	-	0.32	-	-	0.78	-	0.48
DP 1219 B2RF	-	-	3.69	-	-	-	-	-	-	-	-	-	1.28	-	-	6.36	-	3.70
DP 1252 B2RF	11.09	-	-	-	18.15	17.52	-	-	-	2.48	0.03	-	-	1.60	-	0.48	-	3.35
DP 1311 B2RF	-	0.28	-	-	-	-	-	0.80	0.51	0.50	*	-	-	-	-	-	-	0.05
DP 1321 B2RF	3.33	1.51	0.32	-	2.57	0.08	5.20	6.39	-	16.44	2.03	-	9.92	0.21	-	0.26	0.50	1.42
DP 1359 B2RF	-	-	7.59	-	-	-	-	-	-	-	-	-	-	-	-	1.43	-	0.89
DP 1410 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	5.44	-	-	0.76	-	0.57
DP 1441 RF	-	-	-	-	-	*	-	-	-	-	-	-	-	-	-	-	-	*
DP 1454 NR B2RF	-	-	-	-	3.20	1.96	-	-	-	-	-	-	-	0.31	-	-	-	0.30
DP 1518 B2XF	1.58	3.76	-	-	-	0.93	3.81	-	3.24	0.33	2.65	-	1.60	2.37	12.24	0.98	1.49	1.40
DP 1522 B2XF	0.23	3.76	-	2.84	0.99	0.44	2.33	0.31	2.29	3.41	4.79	-	1.92	1.74	16.37	1.01	2.32	1.60
DP 1538 B2XF	0.32	-	-	-	5.72	7.06	-	0.26	-	3.44	3.64	-	-	19.94	-	-	0.73	1.88
DP 1549 B2XF	-	-	-	-	-	0.64	-	-	-	-	-	-	-	-	-	1.15	-	0.74
DP 1553 B2XF	0.38	-	-	3.92	15.30	9.90	-	0.07	-	-	0.21	-	-	10.65	-	0.26	-	1.97
DP 1555 B2RF	-	-	-	3.63	5.85	2.94	-	1.35	-	2.54	0.02	-	-	0.94	-	0.05	-	0.64
DP 1558NR B2RF	0.15	-	-	-	5.90	1.37	-	0.07	-	-	0.03	-	-	0.28	-	-	-	0.26
DP 912 B2RF	6.31	1.60	6.05	-	-	0.90	6.91	2.16	5.53	4.17	1.01	-	26.72	0.08	2.44	0.05	0.42	1.57
Deltapine Total	42.22	13.37	41.06	16.18	86.13	63.44	18.25	37.83	11.57	49.16	18.24	-	64.32	40.49	31.40	21.87	7.75	31.15

Table 1. Estimated percentage of upland cotton planted to specified varieties, by states and United States, crop of 2015

	AL	AR	AZ	CA	FL	GA	KS	LA	MO	MS	NC	NM	OK	SC	TN	TX	VA	US
Dyna-Gro																		
DG 2285 B2RF	0.33	-	-	-	-	-	-	-	-	-	0.28	1.00	-	-	-	0.23	-	0.16
DG 2450 B2RF	-	-	-	-	-	-	-	-	-	-	-	3.33	0.40	-	-	-	-	0.02
DG 2570 B2RF	-	0.08	-	-	-	-	-	0.61	2.24	0.37	1.95	10.00	7.20	0.07	0.70	4.18	-	2.79
DG 2595 B2RF	-	-	-	-	-	-	8.10	-	-	-	-	-	-	-	-	1.69	-	0.98
DG 2610 B2RF	-	-	-	-	-	-	-	-	-	-	*	-	-	0.15	-	-	-	*
DG 3385 B2XF	-	-	-	-	-	-	-	-	4.40	-	2.32	-	-	1.12	2.09	*	-	0.28
DG 2355 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1.08	-	0.62
Dyna-Gro Total	0.33	0.08	-	-	-	-	8.10	0.61	6.64	0.37	4.55	14.33	7.60	1.34	2.78	7.18	-	4.85
Phytogen																		
PHY 339 WRF	2.37	-	-	-	0.82	0.38	4.29	1.61	-	1.05	0.63	0.67	-	0.43	-	1.41	1.06	1.08
PHY 367 WRF	-	0.01	-	-	-	0.04	-	-	-	-	0.14	2.17	-	0.06	-	0.34	0.20	0.22
PHY 375 WRF	7.98	0.22	-	-	-	2.60	2.40	-	0.60	-	1.13	21.50	0.48	0.19	1.49	0.01	2.45	0.89
PHY 499 WRF	10.59	1.40	-	6.46	7.88	11.87	-	32.04	1.03	6.16	12.74	17.50	-	16.48	2.63	3.15	29.63	6.07
PHY 575 WRF	1.43	-	-	3.93	-	3.44	-	-	-	-	-	-	-	0.05	-	-	-	0.54
PHY 725 RF	-	-	-	26.78	-	-	-	-	-	-	-	-	-	-	-	-	-	0.15
PHY 755 WRF	-	-	-	8.25	-	-	-	-	-	-	-	-	-	-	-	-	-	0.05
PHY 333 WRF	19.98	4.65	-	-	-	4.87	4.00	10.30	1.62	4.59	16.65	3.67	0.64	7.50	7.62	3.16	20.21	5.02
PHY 417 WRF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.89	-	0.51
PHY 427 WRF	-	-	-	-	-	0.48	-	0.75	-	-	0.43	-	-	0.14	-	-	0.13	0.10
PHY 495 W3RF	2.03	0.06	-	-	1.76	0.08	-	5.48	-	-	0.75	-	-	0.48	1.04	0.40	0.29	0.48
PHY 599 WRF	-	-	-	-	1.65	0.37	-	-	-	-	0.87	-	-	-	-	-	-	0.11
PHY 222 WRF	-	-	-	-	-	0.17	10.83	-	-	-	-	-	-	-	-	0.02	-	0.06
Phytogen Total	44.38	6.34	-	45.42	12.11	24.30	21.52	50.19	3.25	11.80	33.36	45.50	1.12	25.33	12.78	9.38	53.97	15.26
Seed Source Genetics																		
CT 210	-	-	-	-	-	0.05	-	-	-	-	-	-	-	-	-	-	-	0.01
UA 222	-	-	-	-	-	*	-	-	-	-	0.13	-	-	-	-	*	-	0.01
Seed Source Genetics Total	-	-	-	-	-	0.05	-	-	-	-	0.13	-	-	-	-	*	-	0.02
Miscellaneous																		
MISC	-	-	-	4.15	-	0.32	1.19	0.33	-	-	1.18	-	1.68	0.49	-	1.51	0.71	1.05
Miscellaneous Total	-	-	-	4.15	-	0.32	1.19	0.33	-	-	1.18	-	1.68	0.49	-	1.51	0.71	1.05
Total	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00
Transgenic	100.00%	100.00%	100.00%	95.85%	100.00%	99.63%	98.81%	99.54%	100.00%	100.00%	98.66%	100.00%	98.32%	99.48%	100.00%	98.22%	99.29%	98.78%
Conventional	0.00%	0.00%	0.00%	4.15%	0.00%	0.37%	1.19%	0.46%	0.00%	0.00%	1.34%	0.00%	1.68%	0.52%	0.00%	1.78%	0.71%	1.22%

Table 2. Estimated percentage of upland cotton planted to specified varieties, by Classing Office and United States, crop of 2015

Variety	US	Abilene				Corpus Christi	Dumas			Florence			
	Total	KS	OK	TX	Classing Office	TX	AR	MS	Classing Office	NC	SC	VA	Classing Office
ST 4946 GLB2 Bayer CropScience - Stoneville	10.92%	-	0.08%	7.21%	5.85%	4.35%	51.33%	26.07%	30.08%	25.33%	8.37%	24.07%	19.52%
PHY 499 WRF Phytogen	6.07%	-	-	3.51%	2.84%	21.25%	2.61%	6.68%	6.03%	12.74%	16.48%	29.63%	16.03%
DP 1044 B2RF Deltapine	5.62%	-	17.12%	10.54%	11.56%	6.04%	-	-	-	-	-	-	-
PHY 333 WRF Phytogen	5.02%	4.00%	0.64%	0.24%	0.36%	17.40%	3.62%	4.68%	4.51%	16.65%	7.50%	20.21%	14.03%
FM 2011 GT Bayer CropScience - FiberMax	4.72%	-	-	1.56%	1.26%	-	-	-	-	-	-	-	-
FM 2484 B2F Bayer CropScience - FiberMax	4.30%	-	-	3.11%	2.51%	-	-	-	-	-	-	-	-
ST 4747 GLB2 Bayer CropScience - Stoneville	4.18%	26.02%	2.08%	5.84%	5.45%	0.62%	8.96%	-	1.42%	3.33%	1.50%	3.73%	2.77%
FM 9250 GL Bayer CropScience - FiberMax	3.82%	0.66%	-	0.64%	0.52%	-	-	-	-	-	-	-	-
DP 1219 B2RF Deltapine	3.70%	-	1.28%	19.51%	16.02%	8.30%	-	-	-	-	-	-	-
DP 1252 B2RF Deltapine	3.35%	-	-	-	-	4.78%	-	2.84%	2.39%	0.03%	1.60%	-	0.55%
NG 1511 B2RF Americot	3.22%	1.84%	5.52%	2.38%	2.93%	0.86%	1.81%	4.20%	3.82%	0.75%	-	2.07%	0.66%
FM 1944 GLB2 Bayer CropScience - FiberMax	2.92%	9.08%	2.32%	10.31%	8.88%	10.38%	1.81%	-	0.29%	7.81%	2.79%	5.03%	5.80%
FM 2334 GLT Bayer CropScience - FiberMax	2.82%	-	-	2.29%	1.85%	-	-	-	-	-	-	-	-
DG 2570 B2RF Dyna-Gro	2.79%	-	7.20%	4.53%	4.94%	0.25%	-	-	-	1.95%	0.07%	-	1.09%
DP 1553 B2XF Deltapine	1.97%	-	-	0.93%	0.75%	0.79%	-	-	-	0.21%	10.65%	-	3.67%
DP 1050 B2RF Deltapine	1.92%	-	-	-	-	-	-	-	-	-	0.41%	-	0.14%
DP 1538 B2XF Deltapine	1.88%	-	-	-	-	-	-	3.95%	3.32%	3.64%	19.94%	0.73%	8.72%
DP 1522 B2XF Deltapine	1.60%	2.33%	1.92%	0.52%	0.80%	-	2.08%	3.77%	3.51%	4.79%	1.74%	2.32%	3.48%
DP 1137 B2RF Deltapine	1.59%	-	-	-	-	-	-	3.13%	2.63%	1.58%	1.71%	0.10%	1.44%
DP 912 B2RF Deltapine	1.57%	6.91%	26.72%	0.15%	4.95%	-	6.27%	4.74%	4.99%	1.01%	0.08%	0.42%	0.63%
Epic RF All-Tex	1.53%	-	-	0.16%	0.13%	-	-	-	-	-	-	-	-
DP 1321 B2RF Deltapine	1.42%	5.20%	9.92%	0.44%	2.18%	-	6.03%	16.65%	14.97%	2.03%	0.21%	0.50%	1.24%
DP 1518 B2XF Deltapine	1.40%	3.81%	1.60%	-	0.33%	-	2.08%	0.38%	0.65%	2.65%	2.37%	1.49%	2.42%

Table 2. Estimated percentage of upland cotton planted to specified varieties, by Classing Office and United States, crop of 2015

Variety	US	Lamesa			Lubbock	Macon				Memphis					
	Total	NM	TX	Classing Office	TX	AL	FL	GA	Classing Office	AL	AR	MO	MS	TN	Classing Office
ST 4946 GLB2 Bayer CropScience - Stoneville	10.92%	-	11.82%	11.61%	7.88%	2.87%	-	1.76%	1.82%	5.05%	53.06%	48.41%	63.29%	30.97%	38.94%
PHY 499 WRF Phytogen	6.07%	20.00%	0.66%	1.01%	0.38%	13.34%	7.88%	11.87%	11.85%	5.02%	0.99%	1.03%	3.48%	2.63%	2.19%
DP 1044 B2RF Deltapine	5.62%	-	11.78%	11.57%	6.49%	3.03%	10.55%	-	1.09%	-	-	-	-	-	-
PHY 333 WRF Phytogen	5.02%	-	-	-	2.74%	5.49%	-	4.87%	4.67%	49.40%	5.00%	1.62%	5.18%	7.62%	11.95%
FM 2011 GT Bayer CropScience - FiberMax	4.72%	10.00%	7.98%	8.01%	12.64%	-	-	-	-	-	-	-	-	-	-
FM 2484 B2F Bayer CropScience - FiberMax	4.30%	25.00%	7.36%	7.67%	10.52%	-	-	-	-	-	-	-	-	-	-
ST 4747 GLB2 Bayer CropScience - Stoneville	4.18%	-	0.99%	0.97%	6.25%	-	-	0.95%	0.75%	14.56%	20.20%	14.62%	4.31%	8.63%	14.01%
FM 9250 GL Bayer CropScience - FiberMax	3.82%	-	4.34%	4.26%	11.39%	-	-	-	-	-	-	-	-	-	-
DP 1219 B2RF Deltapine	3.70%	-	3.55%	3.49%	1.84%	-	-	-	-	-	-	-	-	-	-
DP 1252 B2RF Deltapine	3.35%	-	-	-	-	16.55%	18.15%	17.52%	17.41%	-	-	-	-	-	-
NG 1511 B2RF Americot	3.22%	10.00%	6.61%	6.67%	5.56%	-	-	0.54%	0.43%	-	3.23%	1.12%	-	1.04%	1.38%
FM 1944 GLB2 Bayer CropScience - FiberMax	2.92%	-	-	-	0.03%	3.28%	1.76%	2.51%	2.58%	-	1.25%	2.52%	1.49%	1.93%	1.59%
FM 2334 GLT Bayer CropScience - FiberMax	2.82%	5.00%	10.50%	10.40%	4.38%	-	-	-	-	-	-	-	-	-	-
DG 2570 B2RF Dyna-Gro	2.79%	-	8.39%	8.24%	3.16%	-	-	-	-	-	0.10%	2.24%	0.41%	0.70%	0.87%
DP 1553 B2XF Deltapine	1.97%	-	-	-	-	0.57%	15.30%	9.90%	8.83%	-	-	-	-	-	-
DP 1050 B2RF Deltapine	1.92%	-	-	-	-	13.42%	0.84%	11.70%	11.30%	-	-	-	-	-	-
DP 1538 B2XF Deltapine	1.88%	-	-	-	-	0.23%	5.72%	7.06%	5.97%	0.50%	-	-	-	-	0.08%
DP 1522 B2XF Deltapine	1.60%	-	0.99%	0.97%	1.42%	-	0.99%	0.44%	0.41%	0.70%	4.33%	2.29%	1.24%	16.37%	5.93%
DP 1137 B2RF Deltapine	1.59%	-	-	-	-	1.97%	17.06%	7.29%	7.08%	9.32%	-	-	-	-	1.53%
DP 912 B2RF Deltapine	1.57%	-	-	-	0.03%	9.42%	-	0.90%	2.11%	-	0.05%	5.53%	0.41%	2.44%	2.24%
Epic RF All-Tex	1.53%	-	6.82%	6.70%	2.57%	-	-	-	-	-	-	-	-	-	-
DP 1321 B2RF Deltapine	1.42%	-	-	-	0.35%	0.16%	2.57%	0.08%	0.24%	9.75%	-	-	-	-	1.60%
DP 1518 B2XF Deltapine	1.40%	-	0.99%	0.97%	1.56%	-	-	0.93%	0.74%	4.79%	4.33%	3.24%	-	12.24%	5.80%

Table 2. Estimated percentage of upland cotton planted to specified varieties, by Classing Office and United States, crop of 2015

Variety	US	Rayville			Visalia				Classing Office
	Total	LA	MS /1	Classing Office	AZ	CA	NM	TX /1	
ST 4946 GLB2 Bayer CropScience - Stoneville	10.92%	1.72%	-	1.58%	14.91%	1.42%	-	-	8.70%
PHY 499 WRF Phytogen	6.07%	32.04%	-	29.47%	-	6.46%	15.00%	-	3.69%
DP 1044 B2RF Deltapine	5.62%	0.06%	-	0.06%	23.39%	4.72%	-	-	14.41%
PHY 333 WRF Phytogen	5.02%	10.30%	-	9.47%	-	-	7.33%	-	0.84%
FM 2011 GT Bayer CropScience - FiberMax	4.72%	-	-	-	-	-	-	-	-
FM 2484 B2F Bayer CropScience - FiberMax	4.30%	-	-	-	1.80%	4.19%	-	10.00%	2.53%
ST 4747 GLB2 Bayer CropScience - Stoneville	4.18%	-	-	-	-	-	-	-	-
FM 9250 GL Bayer CropScience - FiberMax	3.82%	-	-	-	-	-	-	-	-
DP 1219 B2RF Deltapine	3.70%	-	-	-	3.69%	-	-	-	2.05%
DP 1252 B2RF Deltapine	3.35%	-	-	-	-	-	-	-	-
NG 1511 B2RF Americot	3.22%	0.25%	-	0.23%	0.48%	-	-	-	0.27%
FM 1944 GLB2 Bayer CropScience - FiberMax	2.92%	-	-	-	0.63%	-	-	-	0.35%
FM 2334 GLT Bayer CropScience - FiberMax	2.82%	-	-	-	6.34%	3.22%	10.33%	-	5.68%
DG 2570 B2RF Dyna-Gro	2.79%	0.61%	11.00%	1.44%	-	-	20.00%	-	2.28%
DP 1553 B2XF Deltapine	1.97%	0.07%	-	0.06%	-	3.92%	-	-	1.20%
DP 1050 B2RF Deltapine	1.92%	0.27%	-	0.24%	-	-	-	-	-
DP 1538 B2XF Deltapine	1.88%	0.26%	-	0.24%	-	-	-	-	-
DP 1522 B2XF Deltapine	1.60%	0.31%	-	0.28%	-	2.84%	-	-	0.87%
DP 1137 B2RF Deltapine	1.59%	4.35%	-	4.00%	-	-	-	-	-
DP 912 B2RF Deltapine	1.57%	2.16%	-	1.99%	6.05%	-	-	-	3.36%
Epic RF All-Tex	1.53%	-	-	-	-	-	-	-	-
DP 1321 B2RF Deltapine	1.42%	6.39%	65.00%	11.09%	0.32%	-	-	-	0.18%
DP 1518 B2XF Deltapine	1.40%	-	-	-	-	-	-	-	-

Table 3. Estimated percentage of American Pima cotton acreage planted to specified varieties, by states, crop of 2015

		AZ	CA	NM \1	TX \1	US
Phytogen	PHY 805 RF	62.38%	42.63%			40.62%
Phytogen	PHY 811 RF	18.64%	39.86%			32.76%
Deltapine	DP 358 RF	4.60%	7.94%			9.71%
Deltapine	DP 340	0.00%	0.45%			4.36%
Deltapine	DP 357	5.55%	0.00%			4.17%
Miscellaneous	MISC	4.17%	3.52%			3.70%
Phytogen	PHY 800	0.00%	2.98%			2.18%
Hazera	HA 1432	0.00%	1.92%			1.40%
Phytogen	PHY 830	3.81%	0.43%			0.75%
Hazera	HA 211	0.00%	0.27%			0.19%
Public	Pima S-7	0.85%	0.00%			0.10%
All-Tex	P-203	0.00%	0.00%			0.05%

/1 Individual state data withheld