

Cotton

Varieties Planted

2017 Crop



U. S. DEPARTMENT OF AGRICULTURE
Agricultural Marketing Service - Cotton and Tobacco Program
Memphis, Tennessee
September 28, 2017

mp_cn833

Cotton Varieties Planted, United States 2017 Crop

The Deltapine brand of Upland cottonseed was the most popular planted in the United States for the 2017-2018 season, according to the USDA, Agricultural Marketing Service's Cotton and Tobacco Program. The Americot brand was the second most popular followed by Phytogen, Bayer CropScience FiberMax, All-Tex/Dyna-Gro and Bayer CropScience Stoneville.

Deltapine brand varieties were the most popular planted in 2017, accounting for 35.9 percent of the United States acreage. This brand accounted for 56.1 percent of the acreage planted in the southeastern states (Alabama, Florida, Georgia, North Carolina, South Carolina, and Virginia). It accounted for about 61.7 percent in the south central states (Arkansas, Louisiana, Mississippi, Missouri, and Tennessee), 22.5 percent in the southwestern states (Texas, Oklahoma, and Kansas), and 24.1 percent in the western states (Arizona, California, and New Mexico). Deltapine's most popular varieties were DP 1646 B2XF, DP 1518 B2XF, DP 1522 B2XF, and DP 1538 B2XF, accounting respectively for 14.8, 4.7, 4.4, and 1.8 percent of the U.S. Upland cotton acreage.

Americot brand varieties were the second most popular planted in 2017, accounting for 27.0 percent of the United States acreage. These varieties accounted for 10.8 percent of the acreage planted in the southeastern states, 8.1 percent in the south central states, 38.7 percent in the southwestern states, and 0.5 percent in the western states. The most popular Americot varieties were NG 3406 B2XF, NG 4545 B2XF, NG 3500 XF, and NG 3517 B2XF, accounting respectively for 10.9, 4.3, 4.3, and 2.4 percent of the United States acreage planted to Upland cotton.

Phytogen brand varieties were the third most popular planted in 2017. These varieties accounted for 14.4 percent of the acreage planted. They accounted for 21.3 percent of the acreage planted in the southeastern states, 14.8 percent of the acreage in the south central states, 11.4 percent in the southwestern states and 22.4 percent in the western states. The most popular Phytogen varieties were PHY 444 WRF, PHY 333 WRF, and PHY 490 W3FE accounting respectively for 3.9, 3.4, and 2.1 percent of the United States acreage planted to Upland cotton.

Bayer CropScience FiberMax brand varieties were the fourth most popular planted in 2017, accounting for 9.6 percent of the United States acreage. They accounted for 0.4 percent of the acreage planted in the southeastern states, 0.1 percent of the acreage in the south central states, 13.9 percent in the southwestern states and 43.6 percent in the western states. The most popular Bayer CropScience FiberMax brand varieties were FM 1830GLT, FM 2011GT, and FM 2334GLT, accounting respectively for 2.8, 1.2, and 1.2 percent of the United States acreage planted to Upland cotton.

All-Tex/Dyna-Gro brand varieties were the fifth most popular and accounted for about 7.0 percent of the U.S. acreage planted in 2017. Bayer CropScience Stoneville varieties were the sixth most popular and accounted for about 4.6 percent of the 2017 cotton acreage.

Phytogen was the most popular brand of American Pima varieties planted in 2017. Phytogen variety PHY 881 RF accounted for 42.6 percent of the United States Pima acreage. Phytogen's PHY 841 RF was the second most planted American Pima variety and accounted for 17.4 percent of the U.S. crop. Phytogen's PHY 888 RF was the next most popular variety and accounted for 10.2 percent of the U.S. Pima acreage.

Estimates of the percentage of the various varieties of cotton planted in the United States for 2017 were based on informal surveys made by the Cotton and Tobacco Program Classing Offices. Those surveyed included ginners, seed dealers, extension agents, and other knowledgeable sources.

**Estimated percentage of Upland cotton planted to leading specified varieties, by growth area,
2017 crop**

Variety	US	Southeast	South Central	Southwest	Far West
DP 1646 B2XF Deltapine	14.77%	32.77%	13.50%	9.21%	0.44%
NG 3406 B2XF Americot	10.94%	0.08%	6.39%	16.46%	0.27%
DG 3385 B2XF All-Tex/Dyna-Gro	4.95%	0.14%	6.07%	6.61%	-
DP 1518 B2XF Deltapine	4.65%	1.06%	24.33%	1.11%	0.94%
DP 1522 B2XF Deltapine	4.38%	2.02%	13.73%	3.01%	0.64%
NG 4545 B2XF Americot	4.25%	*	-	7.04%	-
NG 3500 XF Americot	4.25%	-	-	7.04%	-
PHY 444 WRF Phytogen	3.92%	8.89%	3.72%	2.35%	-
PHY 333 WRF Phytogen	3.43%	4.49%	4.38%	2.71%	5.88%
FM 1830GLT Bayer CropScience - FiberMax	2.82%	-	-	3.90%	17.93%
ST 4946GLB2 Bayer CropScience - Stoneville	2.61%	3.89%	4.26%	1.65%	4.47%
NG 3517 B2XF Americot	2.36%	-	0.01%	3.90%	-
NG 5007 B2XF Americot	2.28%	9.00%	-	0.53%	-
PHY 490 W3FE Phytogen	2.07%	1.83%	0.14%	2.72%	-
DP 1538 B2XF Deltapine	1.76%	8.04%	0.03%	0.02%	-
DP 1549 B2XF Deltapine	1.68%	0.01%	0.02%	2.65%	2.99%
PHY 312 WRF Phytogen	1.57%	0.48%	3.72%	1.48%	-
DP 1612 B2XF Deltapine	1.50%	0.12%	0.60%	2.28%	0.06%
DP 1555 B2RF Deltapine	1.31%	1.75%	5.07%	0.06%	4.58%
DP 1044 B2RF Deltapine	1.29%	-	-	1.53%	14.41%

**Estimated percentage of Upland cotton planted to leading specified varieties, by growth area,
2017 crop**

Brand	US	Southeast	South Central	Southwest	Far West
Deltapine	35.87%	56.11%	61.71%	22.52%	24.07%
Americot	26.95%	10.83%	8.06%	38.67%	0.47%
Phytogen	14.36%	21.34%	14.81%	11.39%	22.43%
Bayer CropScience - FiberMax	9.62%	0.40%	0.05%	13.93%	43.60%
All-Tex/Dyna-Gro	7.00%	0.62%	6.20%	9.80%	0.11%
Bayer CropScience - Stoneville	4.57%	8.17%	8.15%	2.31%	5.75%
CROPLAN	0.88%	1.81%	1.00%	0.55%	-
Miscellaneous	0.48%	0.72%	0.02%	0.37%	3.58%
Concho	0.28%	-	-	0.46%	-
Seed Source Genetics	*	-	*	-	-

Table 1. Estimated percentage of upland cotton planted to specified varieties, by states and United States, crop of 2017

	AL	AR	AZ	CA	FL	GA	KS	LA	MO	MS	NC	NM	OK	SC	TN	TX	VA	US
ALL-TEX/DYNA-GRO																		
ATX CONCHO B2XF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.09	-	0.05
ATX EDGE B2RF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.04	-	0.02
ATX EPIC RF	-	-	-	-	-	-	-	-	-	-	*	-	0.06	-	-	0.01	-	0.01
ATX LA122	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.12	-	0.07
DG 2285 B2RF	-	-	-	-	-	-	-	-	-	-	0.04	-	-	-	-	0.51	-	0.29
DG 2355 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	0.12	-	-	-	-	*
DG 2400 RF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.04	-	0.02
DG 2570 B2RF	-	-	0.20	-	-	-	-	-	-	-	-	-	0.12	-	-	0.13	-	0.08
DG 2595 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	0.24	-	-	0.27	-	0.16
DG 2615 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	0.12	-	-	-	-	*
DG 3109 B2XF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.85	-	0.48
DG 3385 B2XF	-	1.90	-	-	-	-	-	0.18	24.98	0.68	0.84	-	10.33	0.26	7.00	6.40	-	4.95
DG 3445 B2XF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.51	-	0.29
DG 3526 B2XF	-	-	-	-	-	-	-	0.90	-	-	1.55	-	-	1.38	-	-	-	0.09
DG 3544 B2XF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.26	-	0.15
DG 3635 B2XF	-	-	-	-	-	-	-	-	-	-	-	-	0.24	0.35	-	0.54	-	0.32
DG 3645 B2XF	-	-	-	-	-	-	-	0.32	-	-	-	-	-	-	-	*	-	0.01
DG 3757 B2XF	-	-	-	-	-	0.06	-	-	-	-	0.05	-	-	0.60	-	-	-	0.02
All-Tex/Dyna-Gro Total	-	1.90	0.20	-	-	0.06	-	1.41	24.98	0.68	2.48	-	11.24	2.58	7.00	9.78	-	7.00
AMERICOT																		
AM UA48	-	-	-	-	-	-	-	-	-	-	0.08	-	-	*	-	0.34	-	0.19
NG 1511 B2RF	-	-	-	-	-	0.03	-	-	0.01	-	0.02	-	-	-	-	0.59	0.34	0.34
NG 3306 B2RF	-	-	-	-	-	-	-	-	0.19	-	-	-	-	-	-	0.99	-	0.56
NG 3405 B2XF	-	0.27	-	-	-	-	-	-	2.20	0.52	-	-	-	0.04	2.39	0.28	0.08	0.32
NG 3406 B2XF	0.04	0.95	0.49	-	-	0.03	46.09	-	18.81	-	0.09	-	12.67	0.27	17.22	16.47	0.54	10.94
NG 3500 XF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	7.60	-	4.25
NG 3517 B2XF	-	0.04	-	-	-	-	-	-	-	-	-	-	-	-	-	4.21	-	2.36
NG 3522 B2XF	0.09	0.04	-	-	-	0.20	-	-	1.62	-	5.06	-	-	1.67	1.85	*	10.89	0.38
NG 3640 XF	-	-	0.22	-	-	-	-	-	-	-	-	-	-	-	-	0.49	-	0.28
NG 3699 B2XF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.09	-	0.05
NG 4012 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.25	-	0.14
NG 4111 RF	-	-	-	-	-	-	-	-	-	-	-	-	0.07	-	-	0.34	-	0.19
NG 4545 B2XF	-	-	-	-	-	*	-	-	-	-	-	-	2.24	0.02	-	7.45	-	4.25
NG 4601 B2XF	-	-	-	-	-	0.10	-	-	-	-	0.71	-	-	2.09	-	0.08	1.42	0.13
NG 4689 B2XF	-	-	0.14	-	-	-	-	-	-	-	-	-	-	-	-	0.53	-	0.30
NG 5007 B2XF	9.70	-	-	-	5.86	10.58	-	-	-	-	2.00	-	-	12.38	-	0.57	3.10	2.28
Americot Total	9.83	1.32	0.86	-	5.86	10.94	46.09	-	22.83	0.52	7.96	-	14.97	16.47	21.46	40.29	16.37	26.95
Bayer CropScience - FiberMax																		
ACALA DAYTONA RF	-	-	-	10.41	-	-	-	-	-	-	-	-	-	-	-	-	-	0.07
FM 1320GL	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.58	-	0.32
FM 1740B2F	-	-	9.67	1.83	-	-	-	-	-	-	-	35.14	-	-	-	0.07	-	0.36
FM 1830GLT	-	-	25.44	12.60	-	-	0.35	-	-	-	-	3.51	-	-	-	4.21	-	2.82
FM 1888GL	-	-	0.34	-	-	-	-	-	-	-	-	5.62	-	-	-	0.02	-	0.04

Table 1. Estimated percentage of upland cotton planted to specified varieties, by states and United States, crop of 2017

	AL	AR	AZ	CA	FL	GA	KS	LA	MO	MS	NC	NM	OK	SC	TN	TX	VA	US
FM 1900GLT	-	-	0.54	-	-	-	-	-	-	-	-	-	-	-	-	0.09	-	0.06
FM 1911GLT	-	-	0.72	2.46	-	-	-	-	-	-	-	14.25	-	-	-	1.76	-	1.08
FM 1944GLB2	1.32	-	-	-	-	0.08	-	-	0.06	0.14	0.68	-	-	0.10	-	1.68	0.73	1.03
FM 1953GLTP	-	-	-	1.10	-	-	-	-	-	-	-	-	0.15	-	-	0.27	-	0.17
FM 2007GLT	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1.49	-	0.83
FM 2011GT	-	-	0.29	-	-	-	-	-	-	-	-	5.62	-	-	-	2.13	-	1.22
FM 2322GL	-	-	-	0.73	-	-	-	-	-	-	-	-	0.15	-	-	0.42	-	0.25
FM 2334GLT	-	-	2.91	6.57	-	-	-	-	-	-	-	-	0.15	-	-	1.90	-	1.15
FM 2484B2F	-	-	0.61	-	-	-	-	-	-	-	-	-	-	-	-	0.20	-	0.12
FM 9250GL	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.18	-	0.10
Bayer CropScience - FiberMax Total	1.32	-	40.50	35.69	-	0.08	0.35	-	0.06	0.14	0.68	64.15	0.46	0.10	-	15.00	0.73	9.62
Bayer CropScience - Stoneville																		
ST 4747GLB2	0.64	-	-	-	-	-	0.17	-	0.73	-	0.03	-	-	-	-	0.34	1.59	0.24
ST 4848GLT	-	-	-	0.46	-	-	-	0.96	-	-	7.74	-	-	0.36	-	0.10	3.28	0.34
ST 4946GLB2	0.51	8.42	6.21	1.57	9.38	3.24	-	2.41	3.75	1.13	10.75	3.51	-	1.25	5.53	1.78	4.80	2.61
ST 4949GLT	0.62	7.01	0.34	3.10	-	-	-	4.87	-	1.68	0.05	-	-	-	-	0.27	0.06	0.62
ST 5020 GLT	-	0.05	-	-	-	-	-	-	0.73	-	2.16	-	-	1.26	-	-	0.81	0.12
ST 5115GLT	-	0.04	-	-	-	0.12	-	-	-	0.18	0.29	-	-	0.26	0.80	-	0.02	0.06
ST 5289GLT	-	-	-	-	-	-	-	0.15	-	-	-	-	-	-	-	-	-	*
ST 5517GLTP	-	-	-	0.55	-	-	-	-	-	-	-	-	-	-	-	0.01	-	0.01
ST 6182GLT	5.47	-	-	-	-	0.93	-	-	-	-	1.90	-	-	0.56	-	*	0.57	0.39
ST 6448GLB2	-	0.04	-	-	-	0.52	-	4.92	-	-	0.86	-	-	0.35	-	0.01	-	0.18
Bayer CropScience - Stoneville Total	7.23	15.57	6.55	5.68	9.38	4.82	0.17	13.31	5.22	2.99	23.78	3.51	-	4.03	6.33	2.49	11.14	4.57
Concho																		
257	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.50	-	0.28
Concho Total	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.50	-	0.28
CROPLAN																		
CROPLAN 3475B2XF	-	-	-	-	-	-	-	-	-	-	0.04	-	0.61	-	0.36	0.03	-	0.05
CROPLAN 3787B2RF	-	-	-	-	3.04	0.03	-	0.54	-	-	-	-	-	-	-	0.01	-	0.04
CROPLAN 3885B2XF	-	0.01	-	-	4.69	2.78	-	0.23	1.47	0.49	0.08	-	-	0.46	2.60	0.51	-	0.79
CROPLAN total	-	0.01	-	-	7.72	2.81	-	0.78	1.47	0.49	0.12	-	0.61	0.46	2.96	0.55	-	0.88
DELTAPINE																		
DP 1044 B2RF	-	-	21.93	2.74	-	-	-	-	-	-	-	9.14	2.41	-	-	1.48	-	1.29
DP 1050 B2RF	6.75	-	-	-	7.04	5.87	-	-	-	-	-	-	-	-	-	-	-	0.98
DP 1133 B2RF	-	-	-	-	-	-	-	1.51	-	-	-	-	-	-	-	-	-	0.03
DP 1137 B2RF	-	-	-	-	3.54	0.08	-	-	-	-	0.16	-	-	-	-	-	-	0.04
DP 1212 B2RF	-	-	-	-	7.03	-	-	-	-	-	-	-	-	-	-	0.03	-	0.07
DP 1219 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1.76	-	0.98
DP 1252 B2RF	-	-	-	-	-	1.04	-	-	-	-	-	-	-	-	-	-	-	0.12
DP 1321 B2RF	-	-	-	-	0.71	-	-	0.16	-	-	0.17	-	-	-	-	-	-	0.01
DP 1441 RF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.05	0.04	0.03
DP 1518 B2XF	5.42	50.18	1.72	-	-	0.08	4.93	1.49	19.04	16.82	0.15	-	12.66	0.47	20.90	0.25	-	4.65
DP 1522 B2XF	0.99	5.44	0.73	0.91	0.31	0.94	10.86	6.19	14.61	16.92	4.46	-	10.27	6.93	23.52	2.42	2.32	4.38
DP 1538 B2XF	5.23	-	-	-	3.39	8.78	-	-	-	0.08	7.01	-	-	14.12	-	0.02	2.55	1.76

Table 1. Estimated percentage of upland cotton planted to specified varieties, by states and United States, crop of 2017

	AL	AR	AZ	CA	FL	GA	KS	LA	MO	MS	NC	NM	OK	SC	TN	TX	VA	US
DP 1549 B2XF	-	-	3.32	-	-	-	-	0.21	-	-	0.04	6.33	0.30	-	-	2.84	-	1.68
DP 1553 B2XF	0.44	-	-	-	-	0.79	-	-	-	-	0.05	-	-	5.76	-	0.17	-	0.32
DP 1555 B2RF	-	-	7.67	1.46	6.68	2.88	-	22.90	-	8.36	-	-	-	-	-	0.06	-	1.31
DP 1558NR B2RF	-	-	-	-	-	0.12	-	-	-	-	-	-	-	-	-	0.11	-	0.08
DP 1612 B2XF	0.67	-	0.11	-	-	-	3.51	-	1.67	0.12	0.03	-	0.21	-	1.62	2.42	-	1.50
DP 1614 B2XF	6.13	3.81	-	-	0.16	-	-	-	4.19	2.72	4.96	-	-	1.19	1.74	0.03	5.06	0.88
DP 1639 B2XF	-	-	-	-	0.05	0.04	-	9.35	-	-	3.49	-	5.71	3.38	-	0.06	5.86	0.64
DP 1646 B2XF	29.53	6.11	0.27	1.10	20.31	40.36	-	13.58	-	34.61	14.83	-	33.14	35.51	-	7.59	7.94	14.77
DP 1725 B2XF	0.12	0.90	-	-	0.03	0.34	-	0.92	-	0.22	1.51	-	-	0.89	0.39	0.02	3.92	0.21
DP 1747NR B2X	-	-	-	-	-	0.05	-	-	-	-	-	-	-	-	-	0.23	-	0.14
DELTAPINE Total	55.28	66.45	35.75	6.21	49.27	61.39	19.30	56.30	39.51	79.86	36.86	15.46	64.68	68.25	48.17	19.55	27.69	35.87
PHYTOGEN																		
PHY 220 W3FE	-	-	-	-	2.34	-	6.53	-	-	-	-	-	-	-	-	0.09	-	0.10
PHY 222 WRF	-	-	-	-	-	-	-	-	-	-	-	-	0.46	-	-	0.03	0.11	0.04
PHY 223 WRF	-	-	-	-	-	-	-	-	-	-	0.01	-	-	-	-	0.07	0.19	0.04
PHY 243 WRF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.10	-	0.06
PHY 300 W3FE	0.06	-	-	-	-	-	1.13	-	-	-	0.20	-	0.91	0.02	-	0.42	0.16	0.29
PHY 312 WRF	1.90	11.54	-	-	-	-	-	5.39	-	0.89	0.89	-	-	-	0.27	1.60	0.60	1.57
PHY 330 W3FE	6.66	*	-	0.55	3.55	-	-	0.32	-	-	1.44	-	-	0.62	-	0.08	1.18	0.40
PHY 333 WRF	4.04	3.12	8.55	1.10	0.92	3.09	-	7.69	3.66	4.73	9.76	4.92	0.07	0.49	4.12	2.92	24.17	3.43
PHY 339 WRF	0.08	-	-	-	-	-	-	-	0.73	-	-	-	-	-	2.39	0.09	-	0.14
PHY 340 W3FE	-	0.01	-	-	0.05	-	-	0.10	-	-	1.09	-	-	0.23	0.26	0.04	2.25	0.09
PHY 380 W3FE	0.12	-	-	-	-	-	-	-	-	-	0.04	-	-	-	0.01	-	-	0.01
PHY 427 WRF	-	-	-	-	4.69	-	-	-	-	-	-	-	-	-	0.01	0.06	-	0.07
PHY 444 WRF	7.49	0.09	-	-	13.64	13.11	-	9.26	-	8.80	0.73	-	-	0.94	0.01	2.54	0.43	3.92
PHY 450 W3FE	-	-	-	-	-	*	-	0.27	-	-	1.21	-	-	0.88	-	0.10	0.42	0.12
PHY 460 W3FE	-	-	-	-	-	-	9.48	-	-	-	-	-	-	-	-	0.10	-	0.10
PHY 470 W3FE	-	-	-	-	-	-	-	-	-	-	2.29	-	-	0.45	-	0.03	-	0.10
PHY 487 WRF	-	-	-	-	-	0.01	-	-	-	-	0.37	-	-	0.03	-	-	-	0.01
PHY 490 W3FE	1.22	*	-	-	0.13	0.38	16.95	0.26	-	-	6.16	-	6.13	4.09	0.64	2.36	5.67	2.07
PHY 495 W3RF	0.35	-	-	-	-	0.10	-	-	-	0.42	2.53	-	-	0.02	2.39	0.06	6.98	0.27
PHY 496 W3RF	-	-	-	-	-	-	-	0.25	-	-	-	-	-	-	-	-	-	*
PHY 499 WRF	-	-	-	-	-	1.93	-	4.48	1.54	0.49	0.93	-	0.46	0.03	3.99	0.75	1.36	0.94
PHY 575 WRF	4.41	-	-	-	0.10	0.13	-	-	-	-	-	-	-	-	-	-	-	0.18
PHY 725 RF	-	-	4.06	8.34	2.34	-	-	-	-	-	-	7.03	-	-	-	-	-	0.17
PHY 764 WRF	-	-	-	39.70	-	-	-	-	-	-	-	-	-	-	-	-	-	0.27
PHYTOGEN Total	26.34	14.76	12.61	49.69	27.77	18.75	34.09	28.02	5.94	15.32	27.63	11.95	8.04	7.80	14.09	11.44	43.51	14.36
SEED SOURCE GENETICS																		
SSG UA 222	-	*	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	*
SEED SOURCE GENETICS Total	-	*	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	*
Miscellaneous																		
MISC	-	-	3.53	2.74	-	1.15	-	0.19	-	-	0.49	4.92	-	0.31	-	0.40	0.56	0.48
Miscellaneous Total	-	-	3.53	2.74	-	1.15	-	0.19	-	-	0.49	4.92	-	0.31	-	0.40	0.56	0.48
Total	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00

Table 2. Estimated percentage of upland cotton planted to specified varieties, by Classing Office and United States, crop of 2017

Variety	US	Abilene				Corpus Christi	Dumas			Florence			
	Total	KS	OK	TX	Classing Office	TX	AR	MS	Classing Office	NC	SC	VA	Classing Office
DP 1646 B2XF Deltapine	14.77%	-	33.14%	10.81%	17.96%	20.39%	7.53%	37.35%	31.38%	14.83%	35.51%	7.94%	21.42%
NG 3406 B2XF Americot	10.94%	46.09%	12.67%	5.35%	9.48%	0.63%	0.99%	-	0.20%	0.09%	0.27%	0.54%	0.20%
DG 3385 B2XF All-Tex/Dyna-Gro	4.95%	-	10.33%	12.88%	11.49%	-	1.96%	-	0.39%	0.84%	0.26%	-	0.54%
DP 1518 B2XF Deltapine	4.65%	4.93%	12.66%	0.56%	4.84%	0.34%	49.33%	14.10%	21.15%	0.15%	0.47%	-	0.25%
DP 1522 B2XF Deltapine	4.38%	10.86%	10.27%	3.16%	5.89%	1.67%	6.23%	18.70%	16.21%	4.46%	6.93%	2.32%	5.10%
NG 4545 B2XF Americot	4.25%	-	2.24%	7.20%	5.22%	-	-	-	-	-	0.02%	-	0.01%
NG 3500 XF Americot	4.25%	-	-	0.13%	0.08%	-	-	-	-	-	-	-	-
PHY 444 WRF Phytogen	3.92%	-	-	6.45%	4.00%	6.16%	0.08%	8.05%	6.46%	0.73%	0.94%	0.43%	0.77%
PHY 333 WRF Phytogen	3.43%	-	0.07%	-	0.02%	15.05%	4.01%	5.73%	5.39%	9.76%	0.49%	24.17%	8.06%
FM 1830GLT Bayer CropScience - FiberMax	2.82%	0.35%	-	6.64%	4.13%	5.66%	-	-	-	-	-	-	-
ST 4946GLB2 Bayer CropScience - Stoneville	2.61%	-	-	0.13%	0.08%	1.38%	7.52%	1.16%	2.43%	10.75%	1.25%	4.80%	6.71%
NG 3517 B2XF Americot	2.36%	-	-	0.12%	0.07%	-	0.04%	-	0.01%	-	-	-	-
NG 5007 B2XF Americot	2.28%	-	-	-	-	2.93%	-	-	-	2.00%	12.38%	3.10%	5.81%
PHY 490 W3FE Phytogen	2.07%	16.95%	6.13%	4.00%	5.25%	3.65%	*	-	-	6.16%	4.09%	5.67%	5.37%
DP 1538 B2XF Deltapine	1.76%	-	-	-	-	0.10%	-	-	-	7.01%	14.12%	2.55%	9.04%
DP 1549 B2XF Deltapine	1.68%	-	0.30%	4.85%	3.11%	0.43%	-	-	-	0.04%	-	-	0.02%
PHY 312 WRF Phytogen	1.57%	-	-	-	-	9.24%	10.88%	1.11%	3.07%	0.89%	-	0.60%	0.54%
DP 1612 B2XF Deltapine	1.50%	3.51%	0.21%	4.88%	3.23%	-	-	-	-	0.03%	-	-	0.02%
DP 1555 B2RF Deltapine	1.31%	-	-	-	-	-	-	9.46%	7.57%	-	-	-	-
DP 1044 B2RF Deltapine	1.29%	-	2.41%	2.40%	2.30%	4.17%	-	-	-	-	-	-	-
FM 2011GT Bayer CropScience - FiberMax	1.22%	-	-	2.71%	1.68%	0.22%	-	-	-	-	-	-	-
FM 2334GLT Bayer CropScience - FiberMax	1.15%	-	0.15%	5.06%	3.19%	-	-	-	-	-	-	-	-
FM 1911GLT Bayer CropScience - FiberMax	1.08%	-	-	0.17%	0.10%	-	-	-	-	-	-	-	-

Table 2. Estimated percentage of upland cotton planted to specified varieties, by Classing Office and United States, crop of 2017

Variety	US	Lamesa	Lubbock	Macon				Memphis					
	Total	TX	TX	AL	FL	GA	Classing Office	AL	AR	MO	MS	TN	Classing Office
DP 1646 B2XF Deltapine	14.77%	11.61%	0.80%	33.77%	20.31%	40.36%	38.15%	19.62%	5.64%	-	21.40%	-	5.53%
NG 3406 B2XF Americot	10.94%	3.88%	30.40%	-	-	0.03%	0.02%	0.14%	0.94%	18.81%	-	17.22%	9.74%
DG 3385 B2XF All-Tex/Dyna-Gro	4.95%	6.10%	7.16%	-	-	-	-	-	1.88%	24.98%	4.00%	7.00%	9.10%
DP 1518 B2XF Deltapine	4.65%	-	0.25%	-	-	0.08%	0.06%	18.05%	50.46%	19.04%	32.60%	20.90%	29.31%
DP 1522 B2XF Deltapine	4.38%	5.21%	1.40%	-	0.31%	0.94%	0.74%	3.30%	5.18%	14.61%	10.47%	23.52%	12.76%
NG 4545 B2XF Americot	4.25%	5.22%	11.16%	-	-	*	-	-	-	-	-	-	-
NG 3500 XF Americot	4.25%	0.42%	15.30%	-	-	-	-	-	-	-	-	-	-
PHY 444 WRF Phytogen	3.92%	3.12%	0.06%	8.70%	13.64%	13.11%	12.34%	4.66%	0.09%	-	8.87%	0.01%	1.27%
PHY 333 WRF Phytogen	3.43%	1.08%	0.09%	1.39%	0.92%	3.09%	2.67%	10.22%	2.82%	3.66%	0.87%	4.12%	4.08%
FM 1830GLT Bayer CropScience - FiberMax	2.82%	8.78%	0.83%	-	-	-	-	-	-	-	-	-	-
ST 4946GLB2 Bayer CropScience - Stoneville	2.61%	3.31%	1.60%	0.69%	9.38%	3.24%	3.10%	0.07%	8.72%	3.75%	1.20%	5.53%	5.00%
NG 3517 B2XF Americot	2.36%	0.68%	8.29%	-	-	-	-	-	0.05%	-	-	-	0.01%
NG 5007 B2XF Americot	2.28%	-	0.15%	13.86%	5.86%	10.58%	10.93%	-	-	-	-	-	-
PHY 490 W3FE Phytogen	2.07%	1.52%	1.87%	1.51%	0.13%	0.38%	0.57%	0.55%	*	-	-	0.64%	0.24%
DP 1538 B2XF Deltapine	1.76%	-	-	3.40%	3.39%	8.78%	7.54%	9.50%	-	-	-	-	1.09%
DP 1549 B2XF Deltapine	1.68%	8.26%	0.99%	-	-	-	-	-	-	-	-	-	-
PHY 312 WRF Phytogen	1.57%	-	0.06%	-	-	-	-	6.33%	11.76%	-	-	0.27%	4.09%
DP 1612 B2XF Deltapine	1.50%	4.13%	1.96%	-	-	-	-	2.22%	-	1.67%	0.73%	1.62%	1.18%
DP 1555 B2RF Deltapine	1.31%	-	-	-	6.68%	2.88%	2.56%	-	-	-	-	-	-
DP 1044 B2RF Deltapine	1.29%	0.84%	0.19%	-	-	-	-	-	-	-	-	-	-
FM 2011GT Bayer CropScience - FiberMax	1.22%	4.45%	1.74%	-	-	-	-	-	-	-	-	-	-
FM 2334GLT Bayer CropScience - FiberMax	1.15%	5.34%	0.09%	-	-	-	-	-	-	-	-	-	-
FM 1911GLT Bayer CropScience - FiberMax	1.08%	3.30%	2.00%	-	-	-	-	-	-	-	-	-	-

Table 2. Estimated percentage of upland cotton planted to specified varieties, by Classing Office and United States, crop of 2017

Variety	US	Rayville			Visalia				Classing Office
	Total	LA	MS	Classing Office	AZ	CA	NM	TX /1	
DP 1646 B2XF Deltapine	14.77%	13.58%	36.65%	15.33%	0.27%	1.10%	-		0.36%
NG 3406 B2XF Americot	10.94%	-	-	-	0.49%	-	-		0.22%
DG 3385 B2XF All-Tex/Dyna-Gro	4.95%	0.18%	-	0.17%	-	-	-		-
DP 1518 B2XF Deltapine	4.65%	1.49%	-	1.37%	1.72%	-	-		0.77%
DP 1522 B2XF Deltapine	4.38%	6.19%	5.97%	6.17%	0.73%	0.91%	-		0.53%
NG 4545 B2XF Americot	4.25%	-	-	-	-	-	-		-
NG 3500 XF Americot	4.25%	-	-	-	-	-	-		-
PHY 444 WRF Phytogen	3.92%	9.26%	28.21%	10.70%	-	-	-		-
PHY 333 WRF Phytogen	3.43%	7.69%	-	7.10%	8.55%	1.10%	4.92%		6.58%
FM 1830GLT Bayer CropScience - FiberMax	2.82%	-	-	-	25.44%	12.60%	3.51%		18.67%
ST 4946GLB2 Bayer CropScience - Stoneville	2.61%	2.41%	-	2.22%	6.21%	1.57%	3.51%		5.20%
NG 3517 B2XF Americot	2.36%	-	-	-	-	-	-		-
NG 5007 B2XF Americot	2.28%	-	-	-	-	-	-		-
PHY 490 W3FE Phytogen	2.07%	0.26%	-	0.24%	-	-	-		-
DP 1538 B2XF Deltapine	1.76%	-	2.74%	0.21%	-	-	-		-
DP 1549 B2XF Deltapine	1.68%	0.21%	-	0.19%	3.32%	-	6.33%		2.45%
PHY 312 WRF Phytogen	1.57%	5.39%	-	4.98%	-	-	-		-
DP 1612 B2XF Deltapine	1.50%	-	-	-	0.11%	-	-		0.05%
DP 1555 B2RF Deltapine	1.31%	22.90%	26.42%	23.17%	7.67%	1.46%	-		4.86%
DP 1044 B2RF Deltapine	1.29%	-	-	-	21.93%	2.74%	9.14%		15.35%
FM 2011GT Bayer CropScience - FiberMax	1.22%	-	-	-	0.29%	-	5.62%		0.99%
FM 2334GLT Bayer CropScience - FiberMax	1.15%	-	-	-	2.91%	6.57%	-		4.95%
FM 1911GLT Bayer CropScience - FiberMax	1.08%	-	-	-	0.72%	2.46%	14.25%		4.79%

/1 Individual state data withheld

Table 3. Estimated percentage of American Pima cotton acreage planted to specified varieties, by states, crop of 2017

		AZ	CA	NM \1	TX	US
Phytogen	PHY 881 RF	27.82%	45.11%		27.13%	42.57%
Phytogen	PHY 841 RF	15.33%	17.50%		19.38%	17.40%
Phytogen	PHY 888 RF	5.01%	9.97%		15.50%	10.18%
Hazera	HA1432	-	11.45%		-	9.87%
Deltapine	DP 348 RF PIMA	33.56%	3.65%		17.05%	6.73%
Phytogen	PHY 811 RF	3.70%	5.58%		0.00%	5.02%
Phytogen	PHY 805 RF	11.92%	3.39%		16.28%	4.81%
Miscellaneous Pima	MISC -PIMA	2.67%	2.03%		4.65%	2.27%
Deltapine	DP 358 RF PIMA	-	1.32%		-	1.14%

/1 Individual state data withheld